

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number  
**WO 00/79267 A2**

- (51) International Patent Classification<sup>7</sup>: G01N 33/48 (74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).
- (21) International Application Number: PCT/GB00/02446
- (22) International Filing Date: 22 June 2000 (22.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
9914589.8 22 June 1999 (22.06.1999) GB  
0008161.2 3 April 2000 (03.04.2000) GB
- (71) Applicant (for all designated States except US): SCHOOL OF PHARMACY, UNIVERSITY OF LONDON [GB/GB]; 29-39 Brunswick Square, London WC1N 1AX (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): NIZETIC, Dean [HR/GB]; School of Pharmacy, University of London, 29-39 Brunswick Square, London WC1N 1AX (GB). GROET, Jürgen [DE/GB]; School of Pharmacy, University of London, 29-39 Brunswick Square, London WC1N 1AX (GB).
- Published:  
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 00/79267 A2

(54) Title: TREATMENT OF CANCER

(57) Abstract: The product of a gene located at human chromosome 21q11-21 has ubiquitin specific protease activity and binds to ubiquitin, polyubiquitin, ubiquitin-like protein SUMO-3, and proteins which are implicated in the repair/excision of DNA. It is believed to have a role in regulating cell growth, cell growth arrest and/or apoptosis and is implicated in carcinoma growth. The gene has been sequenced and cloned into a microorganism and the product's effect on cell growth is to be investigated. The gene has GenBank accession number AF134213 and has the HUGO-approved name USP25.

### Treatment of Cancer

The present invention relates to the use of the product of the USP25 gene, which appears to have ubiquitin specific protease activity and may have protease activity on ubiquitin-like proteins, located at human  
5 chromosome 21q11 in the treatment, prophylaxis or diagnosis of cancer, especially solid tumours, more particularly lung cancer.

Ubiquitin mediated proteolysis by the 26S proteasome is responsible for the physiological regulation of the levels of many proteins which are key cell cycle and cell growth regulators (Hochstrasser 1995), as well as known  
10 tumour suppressors (Lane 1998).

Ubiquitin is a 76 amino acid polypeptide present in all eukaryotic cells, and highly conserved in evolution. Ubiquitin is conjugated to a target protein through an isopeptide bond between the  $\epsilon$ -amino group of Lys in a target protein (ubiquitination), a process mediated by three groups of  
15 enzymes: ubiquitin activating enzymes (E1), ubiquitin conjugating enzymes (E2) and ubiquitin ligases (E3). Ubiquitinated proteins exist in a monoubiquitinated form, or a multiubiquitin chain: the former is not a degradation signal, while the latter, Lys-48-linked ubiquitin-ubiquitin(n) conjugate, works as a strong degradation signal when joined to a Lys in a  
20 target protein. Protein conjugated to polyubiquitin is then very rapidly and efficiently degraded by non-lysosomal, ATP-dependent degradation by the 26-S proteasome. The de-ubiquitinating enzymes (DUB's) broadly fall into two classes; ubiquitin specific proteases (USPs) and ubiquitin-C-terminal hydrolases (UCHs) are each capable of de-conjugating the ubiquitin-  
25 ubiquitin and ubiquitin-protein links, thereby converting polyubiquitin into mono-ubiquitin, and de-coupling (deubiquitinating) ubiquitin from the target protein, with the result of preventing the degradation of the target protein.

The class of UCH enzymes tends to include relatively small proteins (about 35-40kD) which have low specificity for ubiquitinated proteins. BAP1 is  
30 a UCH but is unusual in that it is larger, having a weight of nearly 100kD.

Several such enzymes are known, the sequences of which show some sequence homologies especially in two domains, the Cys and His domains.

Very broadly, two main functions have been observed among the various members of the USP (UBP) superfamily (Wilkinson 1997, Wilkinson and Hochstrasser 1998). The first is the generation of free ubiquitin from precursor fusion proteins or from peptide-linked polyubiquitin after proteolysis of the targeted protein, and the second is de-ubiquitination. When a protein is targeted for ubiquitin mediated degradation, it is linked to ubiquitin via an isopeptide bond between the C-terminus of ubiquitin and a lysine  $\epsilon$ -amino group(s) of the acceptor protein. Once the conjugate is formed, it can have only two fates: non-lysosomal proteolysis mediated by the 26S proteasome resulting in total protein degradation, or de-conjugation from ubiquitin (de-ubiquitination), resulting in the rescue of the target protein from degradation (Wilkinson 1997, Wilkinson and Hochstrasser 1998).

The regulation of p53 levels by USP25, and consequential prescribed increased cellular predisposition to apoptosis (programmed cell death) upon overexpression of USP25, could be a mechanism by which neuronal loss occurs in Alzheimer's Disease associated with Down Syndrome. Some mechanism could be invoked in explaining the sporadic AD if overexpression of USP25 occurs through a somatic change in aging neurons.

The potential USP catalysed de-ubiquitination and rescue from degradation of tumour suppressors such as p53, could perhaps explain the link to deletions of USPs in solid tumours. De-ubiquitination could play a major role in the Mdm2 mediated control of p53 levels and its activation mechanism, since the ubiquitin-mediated proteasome degradation of p53 is an important effector arm of this recently revealed control pathway (Haupt et al., 1997, Kubbutat et al., 1997, Lane 1998). Halving the dose of a USP could give a pre-cancerous cell a selective advantage in proliferation, by diminishing its rate of "re-cycling" of p53, making it more difficult to achieve a threshold concentration of p53, necessary for its activation. The same hypothetical, gene-dose regulated, tumour suppression model could also

explain why trisomy 21 (Down's syndrome DS) seems to confer protection from development of solid tumours. Among DS children, neuroblastomas and nephroblastomas, and among DS adults, gynaecologic, digestive and breast cancers have very rarely been reported, and are significantly under-  
5 represented compared to the age-matched euploid population (Oster et al., 1975, Satge et al., 1998).

Baker *et al.*, 1999 also propose a role for USP's in tumour formation and cell growth.

In recent years a number of other protein modifying polypeptide tags  
10 have been identified. Many of these are related to ubiquitin and have high levels of identity and similarity (determined using the BLAST algorithm, for instance) to ubiquitin itself. There is a recognised super family of such proteins which have been termed ubiquitin-like proteins (UbL) (Gong et al. 1997, Schwarz et al. 1998). The yeast Smt3 and human SUMO-1 (PIC1,  
15 Sentrin, hSmt3C), SUMO-2 (hSmt3A) and SUMO-3 (hSMT3B) belong to the same family of UbL proteins with approximately 50% identity between themselves, and some 15-30% identity and 40-60% similarity in amino acid sequence to ubiquitin (Lapenta et al. 1997, Mannen et al. 1996, Kamitani et al. 1998, Saitoh and Hinchey, 2000). Yeast and human UBC9 are capable of  
20 conjugating equally yeast or human UbL-s, but not ubiquitin (Schwarz et al. 1998). The SUMO-1,-2 and -3 have the C-terminal glycine, necessary for ubiquitination of the target protein's lysine residue, but unlike ubiquitin, do not have the Lys48 residue necessary for the formation of polyubiquitin chains through isopeptide bonds, which are the signal for the proteasome  
25 degradation (Saitoh and Hinchey, 2000). Nevertheless, yeast Smt3 protein can rescue the mutant Mif2 phenotype, a deficient centromere binding protein resulting in chromosome missegregation (Meluh and Koshland 1995). SUMO-1, as well as SUMO-3 (and probably also SUMO-2) are all capable of being attached by UBC9 to RanGap1, a Ran GTP-ase activating  
30 protein (Kamitani et al. 1998). This ATP-dependent attachment is essential for the binding between modified RanGap1 and RanBP2 binding protein, in

order to form functional nuclear pore complex, which controls export and import of molecules through the nuclear envelope (Mahajan et al. 1997, Matunis et al. 1998, Lee et al. 1998). In addition, UbL small proteins have been shown to modify the death domains of Fas (Okura et al. 1996), Tumour  
5 necrosis factor receptor1 (Okura et al. 1996), PML (a tumour suppressor implicated in the pathogenesis of acute promyelocytic leukaemia) (Kamitani et al. 1998b) and Rad51/52 DNA repair proteins (Shen et al. 1996a). Their conjugating enzyme, UBC9, has been shown to interact by Y2H technique with RAD51/52 DNA repair proteins, and the master tumour suppressor p53  
10 (Shen et al. 1996b). Another UbL is NEDD8 (Kamitani *et al* 1997).

UbL's are conjugated and cleared from their targets by enzymes. Several UbL hydrolase enzymes have been identified which convert precursor UbL to active UbL. Some such enzymes interact with ubiquitin itself as well as with other UbL's. Proteases involved in cleavage of  
15 conjugates of UbL with target protein have been identified for instance SENP1 and SUSP-1, which were recently cloned (Kim et al. 2000, Gong et al. 2000a), and found to specifically cleave SUMO-1,-2 and -3, but not ubiquitin and NEDD8. The first human enzyme with classical USP structure (Cys, His domains) for which dual specificity to both ubiquitin and ubiquitin  
20 like protein was demonstrated was very recently published USP21 on chromosome 1q21 (Gong et al. 2000b). However, opposite from SENP1 and SUSP-1, this enzyme cleaves ubiquitin and Nedd8, but not SUMO-1,-2 or -3 (Gong et al. 2000b).

The proximal third of the chromosome 21 long arm is an exceptionally  
25 gene-poor region of the human genome as estimated by a number of criteria (Shimizu et al., 1995, Yaspo et al., 1995, Gardiner 1996) the estimates of gene-density range from one gene in a megabase to one gene in six megabases of genomic DNA. Until recently only three full length genes had been mapped in this region: STCH (a member of the hsp70 family) (Brodsky  
30 et al., 1995), RIP140 (protein functionally interacting with a variety of nuclear receptors such as estrogen receptor), (Cavailles et al., 1995) and ANA (a

member of the Tob/BTG1 family of tumour suppressors), (Kohno et al., 1998). This region is also an example of extremely highly methylated regions in the human genome.

5       Groet et al. (1998) describe a high-resolution bacterial contig map of 3.4 Mb of genomic DNA in human chromosome 21q11-q21, encompassing the region of elevated disomic homozygosity in Down's syndrome - associated abnormal myelopoiesis and leukemia, and which has shown a strong association with Alzheimer's disease (AD). It was suggested that the high resolution bacterial clone overlap map should be the basis for deriving  
10       a more complete transcriptional map of that region of the chromosome. It was hoped that this would lead to an explanation of the chromosome 21q11 linkage in familial early onset AD (FEOAD) families. In particular it was suggested that a modifier gene in that region could act together with the presenilin-1 gene to generate or modify the AD phenotype.

15       Further work by the present inventors has revealed a new gene in the proximal third of chromosome 21 and that the product of this gene has ubiquitin specific protease properties. It is postulated that the gene product and USPs generally may have a role in AD. The work has been published in Groet et al (2000) after the first priority date of the present application.

20       Valero, et al. (1999) published after the first priority date of the present application, have, in parallel identified this gene and pointed out the gene product's sequence homologies to known USP's in the conserved peptide domains previously identified e.g. by d'Andrea et al (1998). They postulate a role in Alzheimer's disease. This protein has the HUGO  
25       approved name USP25.

      According to the present invention there is provided the new use of the product of the USP25 gene located at human chromosome 21q 11-21, a non-human homologue thereof or a functional fragment thereof in the manufacture of a medicament for use in the diagnosis, prophylaxis or  
30       treatment of cancer.

The cancer is usually a solid cancer, most often non-small cell lung carcinoma, or skin cancer.

A functional fragment herein means a protein having ubiquitin specific protease activity or UbL specific protease activity and comprising a portion  
5 with sequence homology with the product of the said USP25 gene.

It is believed that the ubiquitin specific protease activity of the protein having sequence ID 1 is responsible for its implication in the pathogenesis of cancer. USP activity may be determined using the technique described in the Examples below, in which using bacteria cotransformed with the USP  
10 gene and with a reporter gene encoding a fusion protein which is a ubiquitin-conjugated detectable protein. The protein may be an enzyme detectable by direct enzyme reaction, by enzyme-linked immune assay techniques, by autoradio-graphically or by direct staining after gel separation under conditions suitable to separate ubiquitin and cleaved protein from fusion  
15 protein.

From experiments conducted to determine with which proteins the product having sequence ID 1 interacts, we have found that there is interaction with ubiquitin, polyubiquitin and various ubiquitin precursors, as well as HHR23A (Matsutani *et al* 1994, GenBank Accession No D21235).  
20 There is also interaction with other ubiquitin-like proteins and with proteins which are known to interact with ubiquitin-like proteins, such as Sumo-3 (Mannen *et al* 1996, Kamitani *et al* 1996, Saitoh *et al* 2000, GenBank Accession No. NM 006937) and ubiquitin-like-specific conjugating enzyme 9 (Schwarz *et al* 1998, Lee *et al* 1998, GenBank Accession No. U 66867). The  
25 isolated protein may therefore be characterised further by having a positive interaction in a yeast-two-hybrid procedure with one or more, preferably all three, proteins having the sequences of GenBank Accession Nos. D 21235, NM 006937 and U 66867. Ubiquitin-like specific protease activity may be determined using techniques analogous to those used to determine ubiquitin  
30 specific protease activity, by using a substrate which is a fusion protein of the ubiquitin-like protein of interest and a detectable protein, and using the

usual separation and immune based or autoradiographic identification techniques.

The gene may be transcribed and translated within a cell line selected as positive for the native gene or an active (in terms of ubiquitin or UbL specific protease activity) fragment thereof. The gene has preferably been introduced into a microorganism or a cell line in a form in which it can be transcribed and translated and the microorganism or the cell-line, as the case may be, has been cultured under conditions whereby the gene is replicated during cell division, and is transcribed and translated into ubiquitin specific protease and the USP is recovered. Preferably the gene in the microorganism is recombinant DNA derived from the mRNA from cells having the active USP gene. Suitably the gene includes sequence ID No. 7, more preferably sequence ID No. 6.

According to a further aspect of the invention there is provided a new use of a protein product having Cys, QQD and His domains specified in sequence IDs numbers 2, 3 and 4, respectively, in the manufacture of a medicament for use in the diagnosis, prophylaxis or treatment of cancer. The protein has ubiquitin specific protease or ubiquitin-like specific protease activity, and has the three specified domains in the USP or UbLSP active conformation. Preferably the protein has, outside the specified domains, some level of sequence homology with sequence ID 1, for instance at least 20%, preferably at least 50%, identity with that sequence, and a level of similarity of at least 50%, preferably at least 70% or more with that sequence (in each case determined using, for instance the BLAST algorithm). Preferably the protein has sequence I.D.1 or sequence I.D.5. Homologues, such as the corresponding mouse product, described in Valero, *et al* 1999 may be used or sequences which have the above levels of identity and similarity with such a sequence.

According to a preferred embodiment the protein has sequence I.D.1.  
According to another preferred embodiment the protein has sequence I.D.5.



According to a further aspect of the invention there is provided an *in vitro* method in which mammalian cells are cultured in the presence of the product of the USP25 gene located at human chromosome 21q11 or of a homologue thereof, or a functional fragment of said product.

5        According to a further aspect of the invention there is provided an *in vitro* method in which mammalian cells are cultured in the presence of a protein product having Cys, QQD and His domains specified in sequence IDs numbers 2, 3 and 4, respectively. Preferably the protein has, outside the specified domains, some level of sequence homology with sequence ID 1,  
10        for instance at least 20%, preferably at least 50%, identity with that sequence, and a level of similarity of at least 50%, preferably at least 70% or more with that sequence (in each case determined using, for instance the BLAST algorithm). Preferably the protein has sequence I.D. 1 or sequence I.D.5. Homologues, such as the corresponding mouse product, described in  
15        Valero, *et al* 1999 may be used or sequences which have the above levels of identity and similarity with such a sequence.

In these *in vitro* methods the effect of the protein on cell growth, cell growth arrest and/or apoptosis is assessed.

The microorganism containing the specified gene has preferably been  
20        transformed with a vector comprising at least a portion of sequence I.D.6 comprising residues 199 to 3367, optionally including an additional 96 b.p. exon inserted after base 2356, and optionally including 5' and/or 3' untranslated regions (UTR's). The vector preferably comprises transcriptional and translational control sequences. The microorganism may  
25        be a yeast but is most conveniently a bacterium.

A mammalian cell transfected with a vector preferably comprises exons making up a sequence comprising residues 199 to 3367 of sequence I.D.6 and optionally also regulatory genomic sequences which are present in wild-type chromosome which lead to enhanced expression activity.

The sequence from residues 199 to 3367 of sequence ID no 6 is specified as sequence ID7 and consists of start codon (ATG), an open reading frame of 3165 nucleotides, and a stop codon (TAA).

According to a further aspect of the invention there is provided a new  
5 use of DNA including the gene located at human chromosome 21q 11-21 or a fragment thereof encoding a functional USP product in the manufacture of a medicament for use in the prophylaxis or treatment of cancer. The DNA is preferably incorporated in a gene therapy vehicle, and is preferably part of a vector, for instance a plasmid vector, or viral vector, capable of transfecting  
10 cells *in vivo*. The medicament usually includes a pharmaceutically acceptable carrier.

The invention is illustrated further in the accompanying drawings in which:

Figure 1 represents a map of human chromosome 21

15 Figure 2 represents a map showing the location of exon trapped products from the experiments reported below

Figure 3 represents sequence homologies of USPs and

20 Figure 4 represents the results of the experiment illustrating USP properties reported below.

The following specific description describes the work which has been carried out.

### Experimental

We identify a portion of human chromosome 21 homozygously  
25 deleted in non-small cell non carcinoma (NSCLC) for further study. The region contained the DNA marker with the highest NSCLC-associated loss of homozygosity (LOH), reported by Kohno *et al.* We found a shared region of overlap (SRO) for the hemizygous loss in other NSCLC. The current work is to identify genes in the SRO which have a potential role in tumour  
30 suppression.

A total of 42 fresh NSCLC cases have been analyzed from the Croatian Tumour Bank (CTB), an initiative with set rules and criteria for accumulation of fresh clinical tumor specimens for molecular studies (Spaventi *et al.*, 1994). Of these, half (including tumors #47 and #61) were samples that were recently studied for LOH of the *NM23-H1* gene (Bosnar *et al.*, 1997), and the other half were fresh tumors (data obtainable from CTB, which also lists the tumor stage in the TNM system, grade, size and survival data). In each case, the tumors and normal lung tissue specimens (as evaluated by the surgeon) were frozen in liquid nitrogen in the operating room and further stored at -70°C. Genomic DNA was isolated using standard procedures (Sambrook *et al.*, 1989). For each sample, 4µm serial frozen sections were cut, mounted on glass slides, and stained with hematoxylin-eosin (H&E). A pathologist confirmed the histologic type of the tumor and evaluated the percentage of normal cells within the tumor. Only samples with less than 20% non-tumor cells were used in this study.

#### Markers and LOH Analysis

Microsatellite analysis was performed using polymerase chain reaction (PCR) with appropriate primer pairs (sequences and PCR conditions as in Genome Data Base, Johns Hopkins University, Baltimore, MD), where the forward primer only from each pair was 5' fluorescently labeled with Applied Biosystems (ABI; Foster City, CA) Big Dyes™ (6-FAM or HEX). Amplification products were analysed using an ABI 310 Genetic Analyzer. Size standards (GeneScan 350) were mixed with every sample for accurate sizing; the separation of the mixture of denatured fragments was achieved by electrophoresis through a 47 cm capillary (module GS STR POP4 C) for approximately 30 min. Raw data were analyzed using GeneScan and Genotyper software. LOH ratios were calculated exactly as described in the GeneScan Applications manual provided by ABI. For each individual allele's fluorescence level, an average of 3 independent electrophoresis-analysis cycles on the ABI 310 was used for calculation.

### Fluorescence In Situ Hybridization (FISH)

Unstained 4µm-thick paraffin block sections were fixed to glass slides, and a standard pretreatment protocol was followed for formalin-fixed, paraffin-embedded slides. P1-derived artificial chromosome (PAC)DNA was labelled with digoxigenin-11-dUTP (Boehringer Mannheim, Germany). Approximately 0.5 µg of each labelled PAC DNA sample was mixed with 5 µg of Cot1 DNA (Gibco BRL, Gaithersburg, MD), precipitated, denatured, allowed to preanneal, and then applied to a denatured slide and hybridized overnight. Slides were washed and signal detected using anti-digoxigenin-rhodamine, followed by DAPI counterstain. Images were captured using a Zeiss Axioskop microscope equipped with a charge-coupled device (CCD) Photometrics, Tucson, AZ) connected to an Apple Powermac 8100 computer. Images were captured on 3 levels of focus, and each level was examined for signals using SmartCapture software (Vysis, Inc., Chicago, IL). Only nuclei with signals were counted in each level, and the number of signals in each cell was determined. B: FISH using a pool of PACs 90B5, 126N20 and BAC 39112 as a probe on the paraffin embedded sections of the tumour #61. Two signal nuclei are predominant. C: FISH using a pool of PACs 73M5 and 135E14 as a probe on the paraffin embedded sections of the tumour #61. Single signal nuclei are predominant.

### Northern Blot Analysis

The cDNAs were labelled by random priming and hybridized to human multiple tissue Northern blots (Clontech, Palo Alto, CA) containing 2 µg polyA + RNA per lane using the protocol recommended by the manufacturer. The exposure was for 14 hr to Molecular Dynamics (Sunnyvale, CA) Phosphorimager screens. The I.M.A.G.E. Consortium (Lennon *et al.*, 1996) cDNA clone ID 824710 and the Unigene clone A002B43 have been used as labelled probes in separate experiments.

### Cleavage Analysis of

#### Ubiquitin-Met- $\beta$ -Galactosidase Fusion Protein

This analysis was performed essentially as described (Everett *et al.*, 1997). Model fusion protein ubiquitin-Met- $\beta$ -galactosidase in a pACYC184 (Cm<sup>r</sup> replicon) was represented by the plasmid pACYC-Ub-Met- $\beta$ -gal, a kind  
5 gift of R. Everett. Plasmid pRB105 containing a *Saccharomyces cerevisiae* ubiquitin-specific protease UBP2 in an IPTG-inducible pBR322 (Amp<sup>r</sup> replicon) was a kind gift of R. Baker, and was used as a positive control. The new gene *USP25* was cloned from nucleotide position 203 to nucleotide  
10 position 3367 (numbering as in GenBank AF 134213) into *SacI*/*SaI* cloning sites of the IPTG-inducible *Escherichia coli* expression vector pQE30 (Qiagen, Chatsworth, CA). The *E. coli* XL-1 blue cells were transformed using a standard rubidium chloride-heat shock method with the combination of pACYC-Ub-Met- $\beta$ -gal and either pQE30 vector, pQE30-*USP25*, or  
15 pRB105, and each of the 3 cotransformants was selected on medium containing chloramphenicol (42  $\mu$ g/ml) and carbenicillin (75  $\mu$ g/ml). Western blots were prepared by electrotransfer to a nitrocellulose membrane (Schleicher & Schull, Keene, NH). The  $\beta$ -galactosidase-containing bands were detected by an anti- $\beta$ -galactosidase polyclonal rabbit antiserum (a kind  
20 gift of R. Everett) using an enhanced chemiluminescence (ECL) assay kit (rpn 2132; Amersham, Arlington Heights, IL) under conditions recommended by the manufacturer.

#### Identification and cloning of *USP25*

25 Twelve sequenced exon-trapped products, when analysed using BLAST-N against public sequence databases, revealed clusters of overlapping cDNA clones. Sequences of our exon-trapped products matched exactly the sequences of the cDNAs forming contigs with a large open reading frame (ORF). In three cases (see Fig.2): from EST 824710 to  
30 AA209364, from AA307805 to AA081200 and from N92952r to Z45010, our trapped exon sequences served to bridge the gaps in the gene sequence

using PCR and suitable restriction, ligation and chain extension techniques. The combined sequence (sequence ID no.5 and GenBank accession number AF134213) revealed a 199 bp 5'UTR, start codon, an ORF of 3165 nucleotides encoding a protein of 1055 amino acids, a stop codon, a 3'UTR of 435 nucleotides and a polyadenylation signal. The total length (without the polyA) assembled is 3803 nucleotides. On multiple human tissue Northern blots (Fig.3) a band of 4.1 kb is visible in all 16 tissues tested (including the normal human lung tissue) with a varying intensity. It is most prominent in skeletal muscle and testis, and the latter tissue also reveals a prominent shorter hybridising transcript of 1.4 kb. All tissues also show a larger weaker band of 4.9 kb, which could be due to an alternative polyadenylation site.

In the course of this analysis, the whole genomic sequence of the two PACs (73M5 and 135E14) became publicly available by the German Human Genome Sequencing Consortium (EMBL accession numbers AJ010597 and AJ010598). Comparison of the genomic sequence with the overlapping cDNA clones and exon sequences revealed that 12 out of the 24 exons had been exon-trapped (hatched rectangles in Fig.2). It also became apparent that the region immediately preceeding the first exon of the gene, comprises the known chromosome 21 CpG island at D21S382 (also known as LL56 Not I linking clone on the Not I physical map of 21q, Ichikawa et al., 1993).

When the deduced polypeptide sequence was compared to Swissprot and other public databases using BLAST-P, a clear pattern of significant homologies ( $e=10^{-6}$  to  $10^{-29}$ ) to proteins across the evolutionary spectrum of eukaryotes was found (Fig.4): all of these proteins belonged to the superfamily of ubiquitin specific proteases (USP-s) or ubiquitin carboxy-terminal hydrolases (UCH-s) (Baker et al., 1992, Swanson et al., 1996, Everett et al., 1997, Wilkinson 1997, Hansen-Hagge et al., 1998, Jensen et al., 1998, Fujiwara et al., 1998, Wilkinson and Hochstrasser 1998, The C. elegans Sequencing Consortium 1998). The polypeptide sequences were most highly conserved around the three domains (the Cys box, the QQD box and the His box, Fig.4) known to be essential for the main function of these

enzymes: the cleavage of ubiquitin at its carboxy terminus from extension proteins (ubiquitin precursors) and ubiquitinated proteins and protein fragments targetted for the degradation by the 26S proteasome pathway (Wilkinson and Hochstrasser 1998). The Cysteine residue at position 178 and the Histidine residues at positions 599 and 607 (marked with an asterix in Fig.4), which were shown to be an absolute requirement for the function of USP-s and UCH-s (Amerik et al., 1997, Hansen-Hagge et al., 1998, Wilkinson and Hochstrasser 1998) were found in the correct positions in the sequence of the new gene. Since this is a first member of the USP family known to map onto human chromosome 21, we named this protein USP25, for Ubiquitin Specific Protease on Chromosome 21.

**The novel protein (USP25) cleaves ubiquitin from carboxy-terminal fusion proteins**

The ability of USP25 to cleave a model ubiquitin fusion protein substrate was investigated by co-expression in *E. Coli*. The complete coding sequence of USP25 was cloned into a T5-driven, IPTG inducible expression vector (pQE30). The new gene USP 21 was cloned from nucleotide position 203 to nucleotide position 3367 (numbering as in sequence ID no. 2 into *Sac*/*Sal* cloning sites of the IPTG-inducible *E. coli* expression vector. As a positive control, the plasmid pRB105 containing a UBP2 gene encoding a *S. Cerevisiae* ubiquitin specific protease in an IPTG inducible and Amp<sup>R</sup> vector was used. The XL-1 blue strain of *E. Coli* was co-transformed with the plasmid containing a ubiquitin-Met- $\beta$ -galactosidase model fusion protein in an IPTG-inducible and chloramphenicol resistant vector, in addition to either pQE30 vector, pQE30-USP25 or the positive control (pRB105). (each of the 3 co-transformants was selected on medium containing chloramphenicol (42  $\mu$ g/ml) and carbenicillin (75  $\mu$ g/ml). Co-transformants were grown to exponential phase, IPTG induced, and the crude protein extracts from these cultures were analysed by Western blot using an anti  $\beta$ -galactosidase antibody. (The western blots were prepared by electro transfer to a nitro cellulose membrane (Schleicher and Schuel.)). The  $\beta$ -

galactosidase containing bands were detected by an anti- $\beta$  galactosidase polyclonal rabbit anti serum using enhanced-chemiluminescence assay kit (ECL, Amersham rpn2132) under conditions recommended by the manufacturer.

5 As can be seen in Fig.4, the uncleaved Ub-Met- $\beta$ -gal substrate (band labelled with an asterix in Fig.4, lane 4) converts to an 8 kDa shorter band (triangle in Fig.4) in the cells co-transformed with either USP25(lanes 5,6) or the yeast UBP2 expressing plasmid (lanes 9,10). Constitutive expression of USP25 (lane 5) is quite sufficient to cleave to completeness the low levels of  
10 model substrate. The more prominent and highly induced band migrating slightly further in the gel than the de-ubiquitinated cleavage product is the truncated form of  $\beta$ -galactosidase expressed by the XL-1 blue bacteria (compare to lanes 1,2 in Fig.4). This result demonstrates that the novel gene product named USP25 can efficiently function as a de-ubiquitinating  
15 enzyme.

From the homologies in the functional domains and from its ability to hydrolyse the bond between the C-terminal double glycine of ubiquitin and the linking methionine residue (Fig.5), it can be concluded that USP25 is a member of ubiquitin specific proteases.

#### 20 **Determination of Proteins with which USP25 interacts**

Functional analysis of USP25 was performed with the aim of detecting the cellular proteins which interact with the USP25 protein through protein-protein interaction, using Yeast-Two-Hybrid (Y2H) approach. *Saccharomyces Cerevisiae* yeast has well characterised ubiquitin activating, conjugating and ligating enzymatic machinery, capable of ubiquitinating  
25 human proteins (Scheffner et al. 1998). A cDNA library from human brain cloned in "prey" vector, was co-transfected to yeast cells with USP25 cloned in "bait" vector. Since ubiquitin cleaving activity of USP25 was proven (Groet et al. 2000), this technique has a theoretical chance of detecting the natural  
30 cellular substrates for ubiquitin cleavage and de-ubiquitination by USP25. Since the action of ubiquitin cleavage is very rapid (Wilkinson and



Hochstrasser 1998), the cleavage and dissociation from its natural substrates for fully active USP25 could preclude the ability to detect the interaction through Y2H. In addition, the artificial cross-de-ubiquitination of yeast's own proteins by an overexpressed USP25 could theoretically be harmful for the yeast cell and/or for the molecular interactions required for the Y2H. For these reasons we performed Y2H using USP25-C178A, a site directed mutant we recently engineered (the mutation being of the key Cys residue in the Cys region) which abolishes the capacity for cleavage of ubiquitin by USP25, but should not interfere with the binding of USP25 to its natural ubiquitinated substrates, since this residue is conserved between all UCH-s and USP-s so far identified.

Y2H experiment using USP25-C178A cloned in the yeast two hybrid "bait" vector pAS2 (Clontech) co-transformed into yeast cells together with a human adult brain cDNA library in the "prey" vector pACT2 (Clontech). Interacting events were visualized by the activation of transcription of all three reporter genes: Ade2, Mel1 and His3. Interacting "prey" sequences were verified by PCR-sequencing on the ABI310 automated sequencer, using universal vector primers, and analysed by BLAST search on non-redundant genome and transcriptome sequence databases. The accession numbers of the sequences found to be interacting, from the GenBank database are given in the table.

**Table 1. Summary of frequency and identities of specific interacting proteins from human brain with USP25-C178A, detected using Yeast-Two-Hybrid technique**

5	Summary of Results by decreasing frequency of detection of baits	Number of specific independent clones "fished" by Y2H	Accession number
	HHR23A	8 clones	D21235
	SUMO-3	8 clones	NM 006937
	human UBC9	5 clones	U66867
10	polyUbiquitin	4 clones	AB009010
	Ubiquitin	3 clones	X04803
	Ran BP2 protein	1 clone	NM 006267
15	Various ubiquitin-like precursors (1 or 2 clones each):	4 clones	
	Other proteins (1 or 2 clones each)	10 clones	

### Conclusions - Interaction with HHR23A

20 DNA repair plays a key role in prevention of carcinogenesis and mutagenesis. This is potentially of special significance to solid tumours which have exposure to UV and chemical carcinogens as the major risk factor, such as cancers of the skin and lung. HHR23A is a homologue of yeast RAD23 protein (Masutani et al. 1994), involved in DNA excision-repair after UV damage and implicated in spindle pole body duplication and cell cycle progression in yeast (Watkins et al. 1993, Biggins et al. 1996). Human homologues HHR23A and B (Masutani et al. 1994) both belong to a group of proteins which, when mutated, lead to Xeroderma Pigmentosum, a rare autosomal recessive disorder associated with a high incidence of sunlight (UV) induced skin cancers.

30 More importantly, hRAD23A has also been isolated as a primary interacting protein by the same Y2H technique using E6AP as a "bait" (Kumar et al. 1999). E6AP (Human Papilloma Virus E6 associated protein)

functions as one of the two so far detected ubiquitin ligases (attaching ubiquitin and labelling for degradation) for the master tumour suppressor p53 (Scheffner et al. 1993). The p53 and HHR23A are the only two so far proven targets for this ubiquitin ligase (Kumar et al. 1999).. Since USP25 shows  
5 high rate of target preference for HHR23A (see Table 1), and both HHR23A and p53 are ubiquitinated by E6AP, it could mean that they are both de-ubiquitinated by USP25. The fact that Y2H with USP25 did not pick up p53 is understandable, because p53 is expressed in small traces (very low level) in normal tissues, and gets only accumulated and activated following DNA  
10 damage or other stimuli for programmed cell death (apoptosis) (Haupt et al. 1997, Kubbutat et al. 1997, Lane 1998). Further experiments are therefore justified to provoke the p53 response, and monitor the effects of USP25 on p53 levels.

Moreover, lack of functional E6AP accelerates the polyglutamine-  
15 induced neuronal cell death in the mouse model for the neurodegenerative disease Spinocerebellar-ataxia 1 (SCA1) (Cummings et al. 1999). Lack of E6AP gene in a mouse expressing the polyglutamine stretch mutation of SCA1 protein dramatically reduces the presence of ubiquitinated intranuclear neuronal inclusions, but drastically accelerates the neuronal  
20 degeneration and cell death (Cummings et al. 1999). A very similar effect has been observed in Huntington's Disease (HD), where a dominant negative mutant of a ubiquitin conjugating enzyme (UBC3), when co-expressed in cultured neurons with the huntingtin protein bearing the polyglutamine extension (mutation causing HD), drastically reduces the  
25 presence of ubiquitinated intranuclear neuronal inclusions, but drastically accelerates the neuronal degeneration and cell death (Saudou et al. 1998). If USP25 de-ubiquitinates a similar set of target proteins to the ones ubiquitinated by E6AP, then the overexpression of USP25 may lead to similar effects as the inhibition of ubiquitin conjugation by E6AP. It is  
30 therefore justified to examine the effects of overexpression of USP25 (and other USP-s) on neuronal toxicity.

### Interaction with human UBC9 and ubiquitin like proteins (UbL) <sup>2</sup>

In yeast, there are 13 ubiquitin conjugating enzymes (E2 enzymes) (Scheffner et al. 1998). Only two (UBC3, mentioned previously in context with HD, and UBC9) are essential for cell cycle progression. Without UBC3  
5 the cell cycle is arrested at the transition point from G1 to S phase, whereas without UBC9 the cell cycle is arrested at the transition point from G2 to M phase (Scheffner et al. 1998). Yeast UBC9, and its mammalian homologue, e.g. human UBC9 have a special function among all other UBC(E2) enzymes, in that they are specifically not conjugating to target proteins the  
10 molecule of ubiquitin, but rather of Ubiquitin-Like small proteins (UbL) (Gong et al. 1997, Schwarz et al. 1998). The yeast Smt3 and human SUMO-1 (PIC1, Sentrin, hSmt3C), SUMO-2 (hSmt3A) and SUMO-3 (hSMT3B) belong to the same family of UbL proteins with approximately 50% identity between themselves, and some 15-30% identity and 40-60% similarity in  
15 amino acid sequence to ubiquitin (Lapenta et al. 1997, Mannen et al. 1996, Kamitani et al. 1998, Saitoh and Hinchey, 2000). Yeast and human UBC9 are capable of conjugating equally yeast or human UbL-s, but not ubiquitin (Schwarz et al. 1998). The SUMO-1, -2 and -3 have the C-terminal glycine, necessary for ubiquitination of the target protein's lysine residue, but unlike  
20 ubiquitin, do not have the Lys48 residue necessary for the formation of polyubiquitin chains through isopeptide bonds, which are the signal for the proteasome degradation (Saitoh and Hinchey, 2000). Nevertheless, yeast Smt3 protein can rescue the mutant Mif2 phenotype, a deficient centromere binding protein resulting in chromosome missegregation (Meluh and  
25 Koshland 1995). SUMO-1, as well as SUMO-3 (and probably also SUMO-2) are all capable of being attached by UBC9 to RanGap1, a Ran GTP-ase activating protein (Kamitani et al. 1998). This ATP-dependent attachment is essential for the binding between modified RanGap1 and RanBP2 binding protein, in order to form functional nuclear pore complex, which controls  
30 export and import of molecules through the nuclear envelope (Mahajan et al. 1997, Matunis et al. 1998, Lee et al. 1998). In addition, UbL small proteins

have been shown to modify the death domains of Fas (Okura et al. 1996),  
Tumour necrosis factor receptor1 (Okura et al. 1996), PML (a tumour  
suppressor implicated in the pathogenesis of acute promyelocytic  
leukaemia) (Kamitani et al. 1998b) and Rad51/52 DNA repair proteins (Shen  
5 et al. 1996a). Their conjugating enzyme, UBC9, has been shown to interact  
by Y2H technique with RAD51/52 DNA repair proteins, and the master  
tumour suppressor p53 (Shen et al. 1996b).

The USP25 Y2H data show clear pattern of interaction in the UBC9  
pathway. Interaction with UBC9 itself, is to our knowledge the first of the kind  
10 demonstration of a direct protein-protein interaction between a USP and a  
conjugating enzyme. Interaction with RanBP2 and SUMO-3 clearly shows  
that USP25 could be sharing the similar target repertoire as UBC9 (Saitoh et  
al. 1997). USP25 may be removing the SUMO (UbL) molecules attached to  
targets by UBC9. Alternatively, USP25 may be preparing the UbL-s for  
15 attachment by UBC9 to targets, by removing the oligopeptide extensions  
after the C-terminal Gly-Gly group from the UbL-s. The only ubiquitin  
protease found to be essential for yeast cell cycle progression, Ulp1, was  
found to be specific to Smt3 removal, and not to ubiquitin (Li and  
Hochstrasser 1999). This protease also had a completely different sequence  
20 from known USP-s and UCH-s. Its human homologues, SENP1 and SUSP-1,  
were recently cloned (Kim et al. 2000, Gong et al. 2000a), and found to  
specifically cleave SUMO-1,-2 and -3, but not ubiquitin and NEDD8, another  
UbL (Kamitani et al. 1997). The first human enzyme with classical USP  
structure (Cys, His domains) for which dual specificity to both ubiquitin and  
25 ubiquitin like protein was demonstrated was very recently published USP25  
on chromosome 1q21 (Gong et al. 2000b). However, opposite from SENP1  
and SUSP-1, this enzyme cleaves ubiquitin and Nedd8, but not SUMO-1,-2  
or -3 (Gong et al. 2000b).

If USP25 cleaves the same UbL proteins to which it binds by Y2H  
30 (see Table 1), it would appear that it is the first classical USP capable of  
cleaving both ubiquitin and human SUMO family, which maybe very

significant to its function. Moreover, it is possible that USP25 is actually specific for SUMO-3 rather than SUMO-1 or SUMO-2. The distinction between these targets would be the first of the kind, but remains to be further confirmed. The previously observed interaction with hRAD23A (Table 1), as well as with other various precursor proteins containing the ubiquitin-like-domains (Table 1), could be linked also to its affinity to a certain type of ubiquitin like domain.

Finally, and very importantly, SUMO molecules have been shown to attach post-translationally to p53 tumour suppressor itself, by the conjugating enzyme UBC9 (Rodrigues et al. 1999, Gostissa et al. 1999). This modification for SUMO-1 in vitro requires only SUMO-1, the SUMO-1 activating enzyme and UBC9. The authors of Rodrigues et al. therefore conclude that SUMO-1 modification pathway acts as a potential regulator of the p53 response and may represent a novel target for the development of therapeutically useful modulators of the p53 response (Rodrigues et al. 1999).

We would propose that the cleavage of ubiquitin like proteins, and its association with UbC9 could be an alternative pathway explaining USP25's role in cell cycle control, and through it, its role in control of programmed cell death, with direct implications in both tumour suppression and neurodegeneration.

### Figure Legends

**Figure 1.** Identification of the Shared Region of Overlap (S.R.O.) for hemizygous deletions in 21q11-q21 in NSCLC. A: Cytogenetic map, Not I long range physical map (Ishikawa, et al., 1993), YAC contig (Nizetic, et al 1994, Shinizu et al 1995 and Bosch et al 1996), and bacterial contig, (Groet et al 1998) are shown in consecutive horizontal layers, respectively, above the line showing the markers used in the LOH analysis (oval symbols). Markers are named as in Genome DataBase (prefix "D21" omitted). In the column under each marker an "X" (symbolizing LOH), "+"

(an absence of LOH) and "U" (un-informative, homozygous result) for that marker in the set of eleven Croatian Tumour Bank (CTB) tumours, or in individual tumours #47 and #61 are shown. NT=not tested. For comparison with our data, markers used as probes on the genomic Southern blot of the NSCLC cell line, and/or in LOH analysis of fresh tumours in the study by Kohno and co-workers (Kohno *et al* 1998) are indicated above the empty bar symbolizing the homozygous deletion they found. In our data, hatched bars indicate hemizygous deletions, and black filled bars indicate segments showing absence of LOH or deletions. Squared symbols "X" and "+" stand for predominantly single and predominantly double signal, respectively, detected by FISH on interphase nuclei of the paraffin embedded sections of the tumour #61, when PAC clones named and indicated as bold lines in the PAC contig above the markers line, were used as probes.

**Figure 2.** Trapped exons (hatched rectangles) and exons deduced from overlapping sequence analysis (white rectangles) defining the exon-intron structure of the new gene USP25. Top half shows two PACs 73M5 and 135E14, also used as FISH probes in Fig. 1, which were the source of genomic DNA for exon trapping. Exon locations on the PACs are shown with vertical bars, and the 50 kbp scale bar refers to this part. Bottom half consists of overlapping cDNA fragments corresponding to exons above them, drawn in the same scale, (500 bp scale bar is shown). Names of cDNA clones are as in dB-EST and UniGene databases, 824710 is the address of the clone in the IMAGE Consortium collection. The complete cDNA sequence for the whole gene is the new GenBank entry with the accession number AF 134213.

**Figure 3.** Comparison of protein sequences of USP25 to other eukaryotic members of the superfamily of USP-s. The protein BAP-1 is actually from the family of Ubiquitin C-terminal Hydrolases, a distinct sub-family of this superfamily, showing homology only in the single key aminoacids in the Cys and His domains. Two reports show the localisations

of the highly homologous sequences for the HAUSP gene to 3p21 (Kashuba, *et al* 1997) and 16p13 (Robinson, *et al* 1998), respectively.

**Figure 4.** Demonstration of the de-ubiquitinating activity of USP25 on a model ubiquitin fusion protein. Western blot of an SDS-PAGE was detected using an anti- $\beta$ -galactosidase antiserum. Lanes 1,2: the *E. coli* XL-1 blue cells alone (in all cases second line of the pair is +1PTG). Lanes 3,4: same cells co-transfected with the model fusion protein encoding plasmid pACYC-UB-Met- $\beta$ -galactosidase protein, band labelled with an asterix). Lanes 5,6: as lanes 3,4 except pQE30-USP25 (full length USP25 gene cloned in the pQE30 expression vector) was added instead of pQE30. Lanes 7,8: same as lanes 3,4 except pRB105 (yeast de-ubiquitinating enzyme UBP2) was transfected instead of pQE30. Lanes 9,10: over-exposure of lanes 7,8. Note the presence of the 8kDa shorter, de-ubiquitinated Met- $\beta$ -galactosidase (band labelled with a triangle).

### References

- Amerik, A. Yu., Swaminathan, S., Krantz, B.A., Wilkinson, K.D., and Hochstrasser, M. (1997). *In vivo* disassembly of free polyubiquitin chains by yeast Ubp14 modulates rates of protein degradation by the proteasome. *EMBO J.* 16: 4826-4838.
- Baker R.T. Tobias JW, Varshavsky A. 1992. Ubiquitin-specific proteases of *Saccharomyces cerevisiae*. *J. Biol Chem* 267:23364-23375.
- Baker, R.T., Wang, X-W., Woollatt, E., White, J.A. and Sutherlands, G.R. Identification, functional characterisation, and chromosomal localisation of USP15, a novel Human USP related to Unp Oncoprotein, and a systematic nomenclature for hUSP's. *Genomics* 59, 264-274 (1999).
- Biggins S. et al. *J. Cell Biol.* 133:1331-1346 (1996)
- Bosch, A., Guimera, J., Gaw, S., Gardiner, K., Chumakov, I., Patterson, D and Estivill, X. Integration of 30 CA-repeat markers into the cytogenetic, genetic and YAC maps of human chromosome 21. *Eur. J. Hum. Genet.* 4, 135-142 1996.



- Bosnar MH, Pavelic K, Krizanac S, Slobodnjak Z, Pavelic J. 1997.  
Squamous cell lung carcinomas: the role of nm23-H1 gene. *J Mol Med*  
75:609-613.
- 5 Brodsky, G., Otterson, G.A., Parry, B.B., Hart, I., Patterson, D., and  
Kaye, F.J. Localization of STCH to Human Chromosome 21q11.1.  
*Genomics*, 30:627-628, 1995.
- Cavailles, V., Dauvois, S., L'Horset, F., Lopez, G., Hoare, S.,  
Kushner, P.J., and Parker, M.G. Nuclear factor RIP140 nodulates  
transcriptional activation by the estrogen receptor. *EMBO J.*, 14:3741-3751,  
10 1995.
- Cummings C.J. et al., *Neuron* 24:879-892 (1999)
- d' Andrea, A and Pellman, D. Deubiquitinating enzymes: A new class of  
biological regulators. *Genomics* 62, 395-405 1999.
- 15 Everett, R.D., Meredith, M., Orr, A., Cross, A., Kathoria, M., and  
Parkinson, J. A novel ubiquitin-specific protease is dynamically associated  
with the PML nuclear domain and binds to a herpesvirus regulatory protein.  
*EMBO J.*, 16:1519-1530, 1997
- Fero, M.L., Randel, E., Gurley, K.E., Roberts, J.M., and Kemp, C.J.  
The murine gene p27<sup>Kip1</sup> is haplo-insufficient for tumour suppression.  
20 *Nature*, 396:177-180, 1998.
- Fujiwara, T., Saito, A., Suzuki, M., Shinomiya, H., Suzuki, T.,  
Takahashi, E., Tanigami, A., Ichiyama, A., Chung, C.H., Nakamura, Y., and  
Tanaka, K. Identification and chromosomal assignment of USP1, a novel  
gene encoding a human ubiquitin-specific protease. *Genomics*, 54:155-158,  
25 1998.
- Gardiner K. 1996. Base composition and gene distribution: critical  
patterns in mammalian genome organisation. *Trends Genet.* 12:519-524.
- Gong L. et al. *J. Biol. Chem.* 272(45):28198-28201 (1997)
- Gong L. et al. *J. Biol. Chem.* 275(5):3355-3359 (2000a)
- 30 Gong L. et al. *J. Biol. Chem.* 275(12):14212-14216 (2000b)

- Gostissa M, Hengstermann A, Fogal V, Sandy P, Schwarz SE, Scheffner M, Del Sal G (1999) Activation of p53 by conjugation to the ubiquitin-like protein SUMO-1. *EMBO J* 1999 Nov 15;18(22):6462-71.
- Groet, J., Ives, J.H., South A.P., Baptista, P.B., Jones, T.A., Yaspo, M-L., Lehrach, H., Potier, M-C., Van Broeckhoven, C., Nizetic, D. Bacterial contig map of the 21q11 region associated with Alzheimer's disease and abnormal myelopoiesis in Down syndrome. *Genome Res.*, 8:385-398, 1998.
- Groet, J., Ives, J.H., Jones, T.A., Danton, M., Flomen, R.H., Sheer, D., Hrasvcan, R., Pavelic, K., and Nizetic, D. Narrowing of the Region of Allelic Loss in 21q 11-21 in squamous Non-small cell Lung Carcinoma and Cloning of a Novel Ubiquitin-specific Protease Gene from the Deleted Segment. *Genes Chromosomes Cancer* 27:153-161 (2000).
- Hansen-Hagge TE, Janssen JWG, Hameister H, Papa FR, Zechner U, Seriu T, Jauch A, Becke D, Hochstrasser M, Bartram CR. 1998. An evolutionarily conserved gene on human chromosome 5q33-q34, UBH1, encodes a novel deubiquitinating enzyme. *Genomics* 49:411-418.
- Haupt Y, Maya R, Kazaz A, Oren M (1997) Mdm2 promotes the rapid degradation of p53, *Nature* 387:296-299.
- Ichikawa, H., Hosoda, F., Arai, Y., Shimizu, K., Ohira, M., and Ohki, M. A *Not I* restriction map of the entire long arm of human chromosome 21. *Nat. Genet.*, 4:361-366 1993.
- Jensen DE, Proctor M, Marquis ST, Gardner HP, Ha SI, Chodosh LA, Ishov AM, Tommerup N, Vissing H, Sekido Y, Minna J, Borodovsky A, Schultz DC, Wilkinson KD, Maul GG, Barlev N, Berger SL, Predergast GC, Rauscher FJ III. 1998. BAP1: a novel ubiquitin hydrolase which binds to the BRCA1 RING finger and enhances BRCA1 - mediated cell growth suppression. *Oncogene* 16:1097-1112.
- Kamitani T. et al. *J.Biol.Chem.* 272(45):28557-28562 (1997)
- Kamitani T. et al. *J.Biol.Chem.* 273(18):11349-11353 (1998a)
- Kamitani T. et al. *J.Biol.Chem.* 273(6):3117-3120 (1998b)

- Kashuba, V.I., Grizatullin, R.Z, Protopopov AI, Allikmets R, Korolev S., Li J, Boldog R, Tory K, Zabarovska V, Marcsek Z, Sumegi J, Klein G, Zabarovsky ER, Kisselev L. 1997. NotI linking/jumping clones of human chromosome 3: mapping of the TFRC, RAB7 and HAUSP genes to regions rearranged in leukemia and deleted in solid tumours. FEBS Lett 419:181-185.
- Kim I.K. et al. J.Biol.Chem. 275(19):14102-14106 (2000)
- Kohno, T., Kawanishi, M., Matsuda, S., Ichikawa, H., takada, M., Ohki, M., Yammamoto, T., and Yokota. J. Homozygous deletion and frequent allelic loss of the 21q11.1-q21.1 region including the ANA gene in human lung carcinoma. Genes Chromosom. Cancer, 21:236-243, 1998.
- Kubbutat M.H.G. et al. Nature 387:299-303 (1997)
- Kumar S. et al. J.Biol.Chem. 274(26):18785-18792 (1999)
- Lane D. 1998. Awakening angels. Nature 394:616-617.
- Lapenta V. et al. Genomics 40:362-366 (1997)
- Lee G.W. et al. J.Biol.Chem. 273(11):6503-6507 (1998)
- Lennon GG, Auffray C. Polymeropoulos M, Soares MB. 1996. The I.M.A.G.E. Consortium: an integrated molecular analysis of genomes and their expression. Genomics 33:151-152.
- Li S.J. and Hochstrasser M., Nature 398:246-251 (1999)
- Mahajan R. et al. Cell 88:97-107 (1997)
- Mannen H. et al. Biochem.Biophys.Res.Comm. 222:178-180 (1996)
- Masutani C. et al. EMBO J. 13(8):1831-1843 (1994)
- Matunis M.J. et al. J.Cell Biol. 140(3):499-509 (1998)
- Meluh P.B. and Koshland D., Mol. Biol. Cell 6:793-807 (1995)
- Nizetic, D., Gellen L., Hamvas R., Mott R., Grigoriev A., Vatcheva R., Zehetner G., Yaspo M.L., Dutriaux A., Lopes C., Delabar J.M., Van Broeckhoven C., Potier M.C. & Lehrach H.; An integrated YAC overlap and "cosmid-pocket" map of the human chromosome 21. Human Molecular Genetics 3(5):759-770 (1994).
- Okura T. et al. J.Immunol. 157:4277-4281 (1996)

- Oster, J., Mikkelsen, A. Mortality and life-table in Down syndrome. *Acto Paediatr. Scand.*, 64: 322-326, 1975.
- Robinson PA, Lomonte P, Leek P, Markham AF, Everett RD, 1998. Assignment of herpesvirus-associated ubiquitin-specific protease gene HAUSP to human chromosome band 16p 13.3 by in situ hybridisation, 5 *Cytogenet. Cell Genet.* 83:100.
- Rodriguez MS, Desterro JM, Lain S, Midgley CA, Lane DP, Hay RT (1999) SUMO-1 modification activates the transcriptional response of p53. *EMBO J* 1999 Nov 15;18(22):6455-61.
- 10 Saitoh H. et al. *Proc.Natl.Acad.Sci.USA* 94:3736-3741 (1997)
- Saitoh H. and Hinchey J., *J.Biol.Chem.* 275(9)6252-6258 (2000)
- Sambrook, J., Fritsch, E.F., & Mamiatis, T *Molecular Cloning:a laboratory manual* 2nd ed. Coldspring Harbor, N.Y. 1989.
- Satge, D., Sasco, A.J., Geneix, A., and Malet, P. Another reason to 15 look for tumor suppressor genes on chromosome 21. *Genes Chromosomes. Cancer* 21:1, 1998.
- Saudou F. et al. *Cell* 95:55-66 (1998)
- Scheffner M. et al. *Cell* 75:495-505 (1993)
- Scheffner M. et al. The Ubiquitin-Conjugation System. In: J.M. Peters, 20 D. Finley and R. Harris (eds.), *Ubiquitin and the Biology of the Cell*, Plenum Press, New York, (1998).
- Schwarz S.E. et al. *Proc.Natl.Acad.Sci.USA* 95:560-564 (1998)
- Shen Z. et al. *Genomics* 36:271-279 (1996a)
- Shen Z. et al. *Genomics* 36:183-186 (1996b)
- 25 Shimizu, N., Antonarakis, S., VanBroeckhoven, C., Patterson, D., Gardiner, K., Nizetic, D., Creau, N., Delabar, J., Korenberg, J., Reeves, R., Doering, J., Ritter, O., and Cuticchia, J. Report of the fifth international workshop on human chromosome 21 mapping 1994. *Cytogenet. Cell Genet.*, 70:147-182, 1995.
- 30 Spaventi R., Pecur, L., Pavelic, K., Pavelic, Z.P., Spaventi, S. and Sambrook, P.J. Human Tumour Bank in Croatia: as possible model for a

small bank as part of the future European Tumour Bank Network. Eur. J. Cancer 30A, 419-1994.

- Swanson DA, Freund CL, Ploder L, McInnes RR, Valle D. 1996. A ubiquitin C-terminal hydrolase gene on the proximal short arm of the X chromosome: implications for X-linked retinal disorders. Hum. Mol. Genet 5:533-538.

The *C. elegans* Sequencing Consortium, 1998. Genome sequence of the nematode *C. elegans*: a platform for investigating biology. Science 282:2012-2018.

- Valero, R., Martany, G., Gonzalez-Angulo, O., Gonzalez-Gonzalez, G., Puellas, L and Gonzales-Duarte, R. USP25, a Novel Gene Encoding a Deubiquitinating Enzyme, is located in the Gene-Poor Region 21q11.2. Genomics 62: 395-405 (1999).

Watkins J.F. et al. Mol. Cell Biol. 13:7757-7765 (1993)

- Wilkinson, K.D. (1997). Regulation of ubiquitin-dependent processes by deubiquitinating enzymes. FASEB J. 11: 1245-1256.

Wilkinson, K.D., and Hochstrasser, M. Deubiquitinating enzymes. In: J.M. Peters, D. Finley and R. Harris (eds.), Ubiquitin and the Biology of the Cell, Plenum Press, New York, 1998.

- Wolozin, B. et al/ Regulation of Apoptosis by Presenilin 1. Neurobiology of Aging. Vol. 19. (15) p 523-527, 1998.

- Yaspo M-L, Gellen L, Mott R, Korn B, Nizetic D, Poustka A, Lehrach H. 1995. Model for a transcript map of human chromosome 21: isolation of new coding sequences from exon and enriched cDNA libraries. Hum. Mol. Genet. 4:1291-1304.

**CLAIMS**

1. Use of the product of the USP25 gene located at human chromosome 21q 11-21 or a non-human homologue thereof or a functional fragment of the product manufacture of a medicament for use in the  
5 diagnosis, prophylaxis or treatment of cancer.
2. Use according to claim 1 in which the cancer is a solid tumour, for instance non-small cell lung carcinoma or skin cancer.
3. Use according to claim 1 or claim 2 in which the product is obtained from a microorganism or cell line in which the gene as recombinant  
10 DNA has been introduced in a form in which it is replicated upon cell division, and is transcribed and translated.
4. Use according to claim 3 in which the recombinant DNA is incorporated in a vector which comprises at least a portion of sequence I.D.6 comprising residues 199 to 3367, optionally including an additional 96 b.p.  
15 exon inserted after base 2356, and optionally including 5' and/or 3' untranslated regions (UTR's).
5. Use of a protein product having Cys, QQD and His domains specified in sequence IDs numbers 2, 3 and 4, respectively, in the manufacture of a medicament for use in the diagnosis, prophylaxis or  
20 treatment of cancer.
6. Use according to claim 5 in which the protein has sequence I.D. No. 1.
7. Use according to claim 5 in which the protein has sequence I.D. No. 5.
- 25 8. An *in vitro* method in which mammalian cells are cultured in the presence of a protein product having Cys, QQD and His domains specified in sequence IDs numbers 2, 3 and 4, respectively, and the effect of protein on cell growth, cell growth arrest and/or apoptosis is assessed.
9. A method according to claim 8 in which the protein has  
30 sequence I.D.1 or sequence I.D.5.

10. Use of DNA including the gene located at human chromosome 21q 11-21 (USP25) or a fragment thereof encoding a functional USP or UbLSP product in the manufacture of a medicament for use in the prophylaxis or treatment of cancer.

- 5 11. Pharmaceutical composition comprising a gene therapy vehicle and DNA including the gene located at human chromosome 21q 11-21 (USP25) or a fragment thereof encoding a functional USP or UbLSP product in transcribable form.

10

Figure 1






Figure 2






Figure 3



4/4

Figure 4



BEST AVAILABLE COPY

## Sequence Listings

5

SEQ ID 1 protein:

10 MTVEQNVLQQSAAQKHQQTFLNQLREITGINDTQILQQALKDSNGNLELAV  
AFLTAKNAKTPQQEETYYQTALPGNDRIYISVGSQADTNVIDLTGDDKDDL  
QRAIALSLAESNRAFRETGITDEEQAISRVLEASIAENKACLKRTPTTEVWRD  
SRNPYDRKRQDKAPVGLKNVGNTCWFSAVIQSLFNLLFRRLLVNLNYKPPS  
NAQDLPRNQKEHRNLPFMRELRYLFALLVGTRKYVDPSRAVEILKDAFK  
15 SNDSQQQDVSEFTHKLLDWLEDAFQMKAEEETDEEKPKNPMVELFYGRF  
LAVGVLEGKKFENTEMFGQYPLQVNGFKDLHECLEAAMIEGEIESLHSEN  
SGKSGQEHWFTELPPVLTFFELSRFEFNQALGRPEKIHNLKLEFPQVLYLDR  
YMHNRNREITRIKREEIKRLKDYLTVLQQRLERYLSYSGSGPKRFPLVDVLQYA  
LEFASSKPVCTSPVDDIDASSPPSGSIPSQTLPTTEQQGALSSELPSTSPS  
20 SVAAISSRSVIHKPFTQSRIPDLPMPAPRHITTEELSVLESCLHRWRTEIE  
NDTRDLQESISRIHRTIELMYSDKSMIQVPYRLHAVLVHEGQANAGHYWAY  
IFDHRESRWMKYNDIAVTKSSWEELVRDSFGGYRNASAYCLMYINDKAQFL  
IQEEFNKETGQPLVGIELPPDLRDFVEEDNQRFEKELEEWDAQLAQKALQ  
EKLLASQKLRESETSVTTAQAAGDPEYLEQPSRSDFSKHLKEETIQITKASH  
25 EHEDKSPETVLQSAIKLEYARLVKLAQEDTPPETDYRLHHVWYFIQNQAPK  
KIIKTLLEQFGDRNLSFDERCHNIMKVAQAKLEMIKPEEVNLEEYEEWHQD  
YRKFRETTMYLIIGLENFQRESYIDSLFLICAYQNNKELLSKGLYRGHDEELI  
SHYRRECLLKLNEQAALFESGEDREVNGLIIMNEFIVPFLPLLLVDEMEEK  
DILAVEDMRNRWCSYLGQEMEPHLQEKLTDPLKLLDCSMEIKSFHEPPKL  
30 PSYSTHELCECFARIMLSLSRTPADGR

SEQ ID no 2: (polypeptide)

GLKNVGNTCWFSAVIQSL

SEQ ID no 3: (polypeptide)

5 QQDVSEFTHKLLDWLED

SEQ ID no 4: (polypeptide)

YRLHAVLVHEGQANAGHYWAY

10 SEQ ID no 5: (polypeptide)

MTVEQNVLQQSAAQKHQQTFLNQLREITGINDTQILQQALKDSNGNLELA  
VAFLTAKNAKTPQQEETYYQTALPGNDRIYISVGSQADTNVIDLTGDDKD  
DLQRAIALSLAESNRAFRETGITDEEQAISRVLEASIAENKACLKRTPTVW  
RDSRNPYDRKRQDKAPVGLKNVGNTCWFSAVIQSLFNLLFRRLVLNYKP

15 PSNAQDLPRNQKEHRNLPFMRELRYLFALLVGTKRKYVDPSRAVEILKDA  
FKSNDSQQQDVSEFTHKLLDWLEDAFQMKAEEETDEEKPKNPMVELFY  
GRFLAVGVLEGKKFENTEMFGQYPLQVNGFKDLHECLEAAMIEGEIESLH  
SENSGKSGQEHWFTELPPVLTFLSRFEFNQALGRPEKIHNKLEFPQVLY  
LDRYMHRNREITRIKREEIKRLKDYLTVLQQRLERYLSYGSGPKRFPLVDV

20 LQYALEFASSKPVCTSPVDDIDASSPPSGSIPSQTLPTTEQQGALSSELP  
STSPSSVAAISSRSVIHKPFTQSRIPDLPMPAPRHITTEKLSVLESCLHR  
WRTEIENDTRDLQESISRIHRTIELMYSKSMIQVPYRLHAVLVHEGQANA  
GHYWAYIFDHRESRWMKYNDIAVTKSSWEELVRDSFGGYRNASAYCLM  
YINDKAQFLIQEEFNKETGQPLVGIETLPPDLRDFVEEDNQRFEKELEEW

25 DAQLAQKALQEKLKLLASQKLRESESVTTAAAGDPEYLEQPSRSDFSKH  
LKEETIQIITKASHEHEDKSPETVLQSIMMTPNMQGIIMAIGKSRSVYDRCG  
PEAGFFKAIKLEYARLVKLAQEDTPPETDYRLHHVVYFIQNQAPKKIIEKT  
LLEQFGDRNLSFDERCHNIMKVAQAKLEMIKPEEVNLEEYEEWHQDYRK  
FRETMYLIIGLENFQRESYIDSLLFLICAYQNNKELLSKGLYRGHDEELIS  
30 HYRRECLLKLNEQAAELFESGEDREVNNGLIIMNEFIVPFLPLLLVDEMEE  
KDILAVEDMRNRWC SYLGQEMEPHLQEKLTDFLPKLLDCSMEIKSFHEP

## PKLPSYSTHELCEFRARIMLSLSRTPADGR

5 SEQ ID no 6:

ds DNA

&gt;BASE COUNT 1223 a 752 c 841 g 987 t

&gt;ORIGIN

1 acagtcggcg ttccgccgcc tgcgcgggt gcccgcgcac gccggccgcc atcgcttcg  
 10 > 61 cgcctggctg gcgggggcgc tgcctccca gccgtccgc gccgtccct ggagctcggc  
 > 121 ggagcgcggc agccagggcc gccggaggcg cgaggagccg gccgccaccg ccgccgccgc  
 > 181 cgcgcggcc gcgggggcca tgaccgtgga gcagaacgtg ctgcagcaga gcgcggcgca  
 > 241 gaagcaccag cagacgtttt tgaatcaact gagagaaatt acggggatta atgacacca  
 > 301 gatactacag caagccttga aggatagtaa tggaaacttg gaattagcag tggcttct  
 15 > 361 tactgcgaag aatgctaaga cccctcagca ggaggagaca acttactacc aaacagcact  
 > 421 tcttggaat gatagataca tcagtgtggg aagccaagca gatacaaatg tgattgatct  
 > 481 cactggagat gataaagatg atcttcagag agcaattgcc ttgagtttg ccgaatcaaa  
 > 541 cagggcattc agggagactg gaataactga tgaggaacaa gccattagca gatttctga  
 > 601 agccagcata gcagagaata aagcatgttt gaagaggaca cctacagaag ttggaggga  
 20 > 661 ttctgaaac ccttatgata gaaaaagaca ggacaaagct cccgttgggc taaagaatgt  
 > 721 tggcaatact tgttggtta gtctgttat tcagtcatta ttaactctt tgaatttag  
 > 781 aagattagtt ctgaattaca agcctccatc aaatgctcaa gatttcccc gaaacaaaa  
 > 841 ggaacatcgg aatttgcctt ttatgcgtga gctgaggtat ctattgcac ttctgttg  
 > 901 taccaaaagg aagtatgtg atccatcaag agcagttgaa attctaagg atgcttcaa  
 25 > 961 atcaaatgac tcacagcagc aagatgtgag tgagtttaca cacaattat tagattggt  
 > 1021 agaagatgcc ttccaaatga aagctgaaga ggagacggat gaagagaagc caaagaaccc  
 > 1081 catgtagag ttgtctatg gcagattcct ggctgtggga gtactgaag gtaaaaaatt  
 > 1141 tgaaaacact gaaatgttg gtcagtacc acttcaggtc aatgggtca aagatctga  
 > 1201 tgagtgccta gaagctgcaa tgattgaagg agaaattgag tctttacatt cagagaattc  
 30 > 1261 aggaaaaatca ggccaagagc attggtttac tgaattacca cctgtgttaa catttgaatt  
 > 1321 gtcaagattt gaatttaac aggcattggg aagaccagaa aaaattcaca acaaattaga  
 > 1381 atttcccaa gtttatatt tggacagata catgcacaga aacagagaaa taacaagaat  
 > 1441 taagaggga gagatcaaga gactgaaaga ttacctcag gtattacaac aaaggctaga  
 > 1501 aagatattha agctatggtt ccggtcccaa acgattcccc ttgtagatg ttctcagta  
 35 > 1561 tgcatggaa ttgcctcaa gtaaactgt ttgcacttct cctgtgacg atattgacgc  
 > 1621 tagttccca cctagtgtt ccataccatc acagacatta ccaagcaca cagaacaaca  
 > 1681 gggagcccta tcttcagaac tgccaagcac atcacctta tcagtgtcg ccatttcac  
 > 1741 gagatcagta atacacaaac catttactca gtcccgata cctccagatt tgcccatga

> 1801 tccggcacca aggcacataa cggaggaaga acttctgtg ctggaaagtt gttlacatcg  
 > 1861 ctggaggaca gaaatagaaa atgacaccag agatttcag gaaagcatat ccagaatcca  
 > 1921 tcgaacaatt gaattaatgt actctgacaa atctatgata caagttcctt atcgattaca  
 > 1981 tgccgtttta gticacgaag gccaagctaa tgctgggcac tactgggcat atattttga  
 5 > 2041 tcatcgtgaa agcagatgga tgaagtacaa tgatattgct gtgacaaaat catcatggga  
 > 2101 agagctagtg agggactctt ttgggtggtta tagaaatgcc agtgcatact gtttaatgta  
 > 2161 cataaatgat aaggcacagt tcctaataca agaggagttt aataaagaaa ctgggcagcc  
 > 2221 ccttgttggg atagaaacat taccaccgga tttagagat ttgttgagg aagacaacca  
 > 2281 acgatttgaa aaagaactag aagaatggga tgcacaactt gccagaaaag ctttcagga  
 10 > 2341 aaagctttta gcgtctcaga aattgagaga gtcagagact tctgtgacaa cagcacaagc  
 > 2401 agcaggagac ccagaatc tagagcagcc atcaagaagt gatttctcaa agcacttgaa  
 > 2461 agaagaaact attcaataa ttaccaaggc atcacatgag catgaagata aaagtcctga  
 > 2521 aacagttttg cagtcggcaa ttaagttgga atatgcaagg ttggttaagt tggccaaga  
 > 2581 agacacccca ccagaaaccg attatcgttt acatcatgta gtggttact ttatccagaa  
 15 > 2641 ccaggcacca aagaaaatta ttgagaaaac attactagaa caatttggag atagaaattt  
 > 2701 gagttttgat gaaaggtgtc acaacataat gaaagttgct caagccaaac tggaaatgat  
 > 2761 aaaacctgaa gaagtaaact tggaggaata tggaggatgg calcaggatt ataggaatt  
 > 2821 cagggaaca actatgtatc tcataattgg gctagaaaat ttcaaagag aaagtatat  
 > 2881 agattccttg ctgttcctca tctgtgctta tcagaataac aaagaactct tgtctaagg  
 20 > 2941 cttatacaga ggacatgatg aagaattgat atcacattat agaagagaat gtttgctaaa  
 > 3001 attaatgag caagccgcag aactcttcga atctggagag gatcgagaag taaacaatgg  
 > 3061 ttgattatc atgaatgagt ttattgtccc attttgcca ttattactgg tggatgaaat  
 > 3121 ggaagaaaag gatatactag ctgtagaaga tatgagaaat cgatgggtgtt cctaccttg  
 > 3181 tcaagaaatg gaaccacacc tccaagaaaa gctgacagat ttttgccaa aactgctga  
 25 > 3241 ttgttctatg gagattaaaa gttccatga gccaccgaag ttaccttcat attccacgca  
 > 3301 tgaactctgt gagcgatttg cccgaalcat gttgtccctc agtcgaactc ctgctgatgg  
 > 3361 aagataaact gcacacttc cctgaacaca ctgtataaac tcttttagt tctaaccct  
 > 3421 tgccttctg tcacaggtt tgctgtgtgc tgctatagt tttaactttt tttatttta  
 > 3481 ataactgcaa aagacaaaal gactatacag actttagtca gactgcagac aataaagctg  
 30 > 3541 aaaatcgcat ggcgtcaga catltaacc ggaactgatg tataatcaca aatctaattg  
 > 3601 attttattat ggcaaaaacta tgctttgcc accttctgt tgcatatta ctttgcttt  
 > 3661 atctttctt tctcaacagc ttccattca gtcggatcc ttcatgact acagccattt  
 > 3721 aaggttcag cactgtglac gatacataat atttggtagc ttgtaaatga aataaagaat  
 > 3781 aaagttttat ttatggctac cta  
 35

SEQ ID no 7 dsDNA

> ca tgaccgtgga gcagaacgtg ctgcagcaga gcgcggcgca  
 > 241 gaagcaccag cagacgtttt tgaatcaact gagagaaatt acggggatta atgacacca

- > 301 gatactacag caagccttga aggatagtaa tggaaacttg gaattagcag tggcttctct
- > 361 tactgcgaag aatgctaaga cccctcagca ggaggagaca acttactacc aaacagcact
- > 421 tcttggcaat gatagataca tcagtgtggg aagccaagca gatacaaatg tgattgatct
- > 481 cactggagat gataaagatg atcttcagag agcaattgcc ttgagtttgg ccgaatcaaa
- 5 > 541 cagggcattc agggagactg gaataactga tgaggaacaa gccattagca gagtctctga
- > 601 agccagcata gcagagaata aagcatgttt gaagaggaca cctacagaag ttggaggga
- > 661 ttctcgaac ccitaitgata gaaaaagaca ggacaaagct cccgttgggc taaagaatgt
- > 721 tggcaatact tgttggttta gtgctgttat tcagtcatta tttaatcttt tggattttag
- > 781 aagattagtt ctgaattaca agcctccatc aaatgctcaa gatttcccc gaaacaaaaa
- 10 > 841 ggaacatcgg aatttgcctt ttatgcgtga gctgaggtat ctatttgcac ttctgttgg
- > 901 taccaaaagg aagtatgttg atccatcaag agcagttgaa attcttaagg atgctttcaa
- > 961 atcaaatgac tcacagcagc aagatgtgag tgagtttaca cacaattat tagattgggt
- > 1021 agaagatgcc ttccaaatga aagctgaaga ggagacggat gaagagaagc caaagaaccc
- > 1081 catggttagag ttgttctatg gcagattcct ggctgtggga gtacttgaag gtaaaaaatt
- 15 > 1141 tgaaaacact gaaatgttg gtcagtaccc acttcaggtc aatgggttca aagatctgca
- > 1201 tgagtgccta gaagctgcaa tgattgaagg agaaattgag tctttacatt cagagaattc
- > 1261 aggaaaatca ggccaagagc attggtttac tgaattacca cctgtgttaa catttgaatt
- > 1321 gtcaagattt gaattaatc aggcattggg aagaccagaa aaaattcaca acaattaga
- > 1381 atttcccaa gttttatatt tggacagata catgcacaga aacagagaaa taacaagaat
- 20 > 1441 taagaggga gagatcaaga gactgaaaga ttacctcag gtattacaac aaaggctaga
- > 1501 aagatattta agctatggtt ccggtcccaa acgattcccc ttggtagatg ttcttcagta
- > 1561 tgcattggaa ttgcctcaa gtaaacctgt ttgcacttct cctgttgacg atattgacgc
- > 1621 tagttccca ctagtggtt ccataccatc acagacatta ccaagcaca cagaacaaca
- > 1681 gggagcccta tctcagaac tgccaagcac atcaccttca tcagtgtctg ccatttcac
- 25 > 1741 gagatcagta atacacaaac catttactca gtcccgata cctccagatt tggccatgca
- > 1801 tccggcacca aggcacataa cggaggaaga actttctgtg ctggaaagt gtttacctg
- > 1861 ctggaggaca gaaatagaaa atgacaccag agatttgag gaaagcatat ccagaatcca
- > 1921 tcgaacaatt gaattaatgt actctgacaa atctatgata caagttcctt atcgattaca
- > 1981 tgcggtttta gttcagaag gccaagctaa tgctgggcac tactgggcat atattttga
- 30 > 2041 tcatcgtgaa agcagatgga tgaagtacaa tgatattgct gtgacaaaat catcatggga
- > 2101 agagctagt agggactctt ttggtgttta tagaatgcc agtgcatact gtttaattga
- > 2161 cataaatgat aaggcacagt tctaataca agaggagttt aataaagaaa ctgggcagcc
- > 2221 ccttgttgt atagaaacat taccaccgga ttgagagat ttgttgagg aagacaacca
- > 2281 acgatttgaa aaagaactag aagaatggga tgcacaactt gccagaaag cttgcagga
- 35 > 2341 aaagctttta gcgtctcaga aattgagaga gtcagagact tctgtgacaa cagcacaagc
- > 2401 agcaggagac ccagaatata tagagcagcc atcaagaagt gattttcaa agcacttgaa
- > 2461 agaagaaact attcaataa ttaccaaggc atcatatgag catgaagata aaagtctga
- > 2521 aacagttttg cagtgcgcaa ttaagttgga atatgcaagg ttggttaagt tggccaaga
- > 2581 agacacccca ccagaacccg attatcgttt acatcatgta gtggtctact ttatccagaa



- > 2641 ccaggcacca aagaaaatta ttgagaaaac attactagaa caatttggag atagaaattt  
> 2701 gagttttgat gaaaggtgtc acaacataat gaaagttgct caagccaaac tggaaatgat  
> 2761 aaaacctgaa gaagttaaact tggaggaata tgaggagtgg catcaggatt ataggaaatt  
> 2821 cagggaaaca actatgtatc tcataattgg gctagaaaat ttcaaagag aaagtatat  
5 > 2881 agattccttg ctgttcctca tctgtgctta tcagaataac aaagaactct tgtctaaagg  
> 2941 cttatacaga ggacatgatg aagaattgat atcacattat agaagagaat gtttgctaaa  
> 3001 attaaatgag caagccgcag aactcttcga atctggagag gatcgagaag taaacaatgg  
> 3061 ttigattatc atgaatgagt ttattgtccc attttgcca ttattactgg tggatgaaat  
> 3121 ggaagaaaag gatatactag ctgtagaaga tatgagaaat cgaagggtgt cctaccttgg  
10 > 3181 tcaagaaatg gaaccacacc tccaagaaaa gctgacagat ttttgccaa aactgcctga  
> 3241 ttgttctatg gagattaaaa gttccatga gccaccgaag ttacctcat attccacgca  
> 3301 tgaactctgt gagcgatttg cccgaatcat gttgtccctc agtcgaactc ctgctgatgg  
> 3361 aagataa

15

## SEQUENCE LISTING

&lt;110&gt; School Of Pharmacy

&lt;120&gt; Treatment of Cancer

&lt;130&gt; HMJ03303WO

&lt;140&gt; PCT/GB 0002446

&lt;141&gt; 2000-06-22

&lt;150&gt; 9914589.6

&lt;151&gt; 1999-06-22

&lt;150&gt; 0008161.2

&lt;151&gt; 2000-04-03

&lt;160&gt; 7

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 1055

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 1

Met Thr Val Glu Gln Asn Val Leu Gln Gln Ser Ala Ala Gln Lys His  
 1 5 10 15

Gln Gln Thr Phe Leu Asn Gln Leu Arg Glu Ile Thr Gly Ile Asn Asp  
 20 25 30

Thr Gln Ile Leu Gln Gln Ala Leu Lys Asp Ser Asn Gly Asn Leu Glu  
 35 40 45

Leu Ala Val Ala Phe Leu Thr Ala Lys Asn Ala Lys Thr Pro Gln Gln  
 50 55 60

Glu Glu Thr Thr Tyr Tyr Gln Thr Ala Leu Pro Gly Asn Asp Arg Tyr  
 65 70 75 80

Ile Ser Val Gly Ser Gln Ala Asp Thr Asn Val Ile Asp Leu Thr Gly  
 85 90 95

Asp Asp Lys Asp Asp Leu Gln Arg Ala Ile Ala Leu Ser Leu Ala Glu  
 100 105 110

Ser Asn Arg Ala Phe Arg Glu Thr Gly Ile Thr Asp Glu Glu Gln Ala  
 115 120 125

Ile Ser Arg Val Leu Glu Ala Ser Ile Ala Glu Asn Lys Ala Cys Leu  
 130 135 140

Lys Arg Thr Pro Thr Glu Val Trp Arg Asp Ser Arg Asn Pro Tyr Asp  
 145 150 155 160

Arg Lys Arg Gln Asp Lys Ala Pro Val Gly Leu Lys Asn Val Gly Asn  
 165 170 175

Thr Cys Trp Phe Ser Ala Val Ile Gln Ser Leu Phe Asn Leu Leu Glu  
 180 185 190

Phe Arg Arg Leu Val Leu Asn Tyr Lys Pro Pro Ser Asn Ala Gln Asp  
 195 200 205

Leu Pro Arg Asn Gln Lys Glu His Arg Asn Leu Pro Phe Met Arg Glu  
 210 215 220

Leu Arg Tyr Leu Phe Ala Leu Leu Val Gly Thr Lys Arg Lys Tyr Val  
 225 230 235 240

Asp Pro Ser Arg Ala Val Glu Ile Leu Lys Asp Ala Phe Lys Ser Asn  
 245 250 255

Asp Ser Gln Gln Gln Asp Val Ser Glu Phe Thr His Lys Leu Leu Asp  
 260 265 270

Trp Leu Glu Asp Ala Phe Gln Met Lys Ala Glu Glu Glu Thr Asp Glu  
 275 280 285

Glu Lys Pro Lys Asn Pro Met Val Glu Leu Phe Tyr Gly Arg Phe Leu  
 290 295 300

Ala Val Gly Val Leu Glu Gly Lys Lys Phe Glu Asn Thr Glu Met Phe  
 305 310 315 320

Gly Gln Tyr Pro Leu Gln Val Asn Gly Phe Lys Asp Leu His Glu Cys  
 325 330 335

Leu Glu Ala Ala Met Ile Glu Gly Glu Ile Glu Ser Leu His Ser Glu  
 340 345 350

Asn Ser Gly Lys Ser Gly Gln Glu His Trp Phe Thr Glu Leu Pro Pro  
 355 360 365

Val Leu Thr Phe Glu Leu Ser Arg Phe Glu Phe Asn Gln Ala Leu Gly  
 370 375 380

Arg Pro Glu Lys Ile His Asn Lys Leu Glu Phe Pro Gln Val Leu Tyr  
 385 390 395 400

Leu Asp Arg Tyr Met His Arg Asn Arg Glu Ile Thr Arg Ile Lys Arg  
 405 410 415

Glu Glu Ile Lys Arg Leu Lys Asp Tyr Leu Thr Val Leu Gln Gln Arg  
 420 425 430

Leu Glu Arg Tyr Leu Ser Tyr Gly Ser Gly Pro Lys Arg Phe Pro Leu  
 435 440 445

Val Asp Val Leu Gln Tyr Ala Leu Glu Phe Ala Ser Ser Lys Pro Val  
 450 455 460

Cys Thr Ser Pro Val Asp Asp Ile Asp Ala Ser Ser Pro Pro Ser Gly  
 465 470 475 480

Ser Ile Pro Ser Gln Thr Leu Pro Ser Thr Thr Glu Gln Gln Gly Ala  
 485 490 495

Leu Ser Ser Glu Leu Pro Ser Thr Ser Pro Ser Ser Val Ala Ala Ile  
 500 505 510

Ser Ser Arg Ser Val Ile His Lys Pro Phe Thr Gln Ser Arg Ile Pro  
 515 520 525

Pro Asp Leu Pro Met His Pro Ala Pro Arg His Ile Thr Glu Glu Glu  
 530 535 540

Leu Ser Val Leu Glu Ser Cys Leu His Arg Trp Arg Thr Glu Ile Glu  
 545 550 555 560

Asn Asp Thr Arg Asp Leu Gln Glu Ser Ile Ser Arg Ile His Arg Thr  
 565 570 575

Ile Glu Leu Met Tyr Ser Asp Lys Ser Met Ile Gln Val Pro Tyr Arg  
 580 585 590

Leu His Ala Val Leu Val His Glu Gly Gln Ala Asn Ala Gly His Tyr  
 595 600 605

Trp Ala Tyr Ile Phe Asp His Arg Glu Ser Arg Trp Met Lys Tyr Asn  
 610 615 620

Asp Ile Ala Val Thr Lys Ser Ser Trp Glu Glu Leu Val Arg Asp Ser  
 625 630 635 640

Phe Gly Gly Tyr Arg Asn Ala Ser Ala Tyr Cys Leu Met Tyr Ile Asn  
 645 650 655

Asp Lys Ala Gln Phe Leu Ile Gln Glu Glu Phe Asn Lys Glu Thr Gly  
 660 665 670

Gln Pro Leu Val Gly Ile Glu Thr Leu Pro Pro Asp Leu Arg Asp Phe  
 675 680 685

Val Glu Glu Asp Asn Gln Arg Phe Glu Lys Glu Leu Glu Glu Trp Asp  
 690 695 700

Ala Gln Leu Ala Gln Lys Ala Leu Gln Glu Lys Leu Leu Ala Ser Gln  
 705 710 715 720

Lys Leu Arg Glu Ser Glu Thr Ser Val Thr Thr Ala Gln Ala Ala Gly  
 725 730 735

Asp Pro Glu Tyr Leu Glu Gln Pro Ser Arg Ser Asp Phe Ser Lys His  
 740 745 750

Leu Lys Glu Glu Thr Ile Gln Ile Ile Thr Lys Ala Ser His Glu His  
 755 760 765

Glu Asp Lys Ser Pro Glu Thr Val Leu Gln Ser Ala Ile Lys Leu Glu  
 770 775 780

Tyr Ala Arg Leu Val Lys Leu Ala Gln Glu Asp Thr Pro Pro Glu Thr  
 785 790 795 800

Asp Tyr Arg Leu His His Val Val Val Tyr Phe Ile Gln Asn Gln Ala  
 805 810 815

Pro Lys Lys Ile Ile Glu Lys Thr Leu Leu Glu Gln Phe Gly Asp Arg  
 820 825 830

Asn Leu Ser Phe Asp Glu Arg Cys His Asn Ile Met Lys Val Ala Gln  
 835 840 845

Ala Lys Leu Glu Met Ile Lys Pro Glu Glu Val Asn Leu Glu Glu Tyr  
 850 855 860

Glu Glu Trp His Gln Asp Tyr Arg Lys Phe Arg Glu Thr Thr Met Tyr  
 865 870 875 880

Leu Ile Ile Gly Leu Glu Asn Phe Gln Arg Glu Ser Tyr Ile Asp Ser  
 885 890 895  
 Leu Leu Phe Leu Ile Cys Ala Tyr Gln Asn Asn Lys Glu Leu Leu Ser  
 900 905 910  
 Lys Gly Leu Tyr Arg Gly His Asp Glu Glu Leu Ile Ser His Tyr Arg  
 915 920 925  
 Arg Glu Cys Leu Leu Lys Leu Asn Glu Gln Ala Ala Glu Leu Phe Glu  
 930 935 940  
 Ser Gly Glu Asp Arg Glu Val Asn Asn Gly Leu Ile Ile Met Asn Glu  
 945 950 955 960  
 Phe Ile Val Pro Phe Leu Pro Leu Leu Leu Val Asp Glu Met Glu Glu  
 965 970 975  
 Lys Asp Ile Leu Ala Val Glu Asp Met Arg Asn Arg Trp Cys Ser Tyr  
 980 985 990  
 Leu Gly Gln Glu Met Glu Pro His Leu Gln Glu Lys Leu Thr Asp Phe  
 995 1000 1005  
 Leu Pro Lys Leu Leu Asp Cys Ser Met Glu Ile Lys Ser Phe His Glu  
 1010 1015 1020  
 Pro Pro Lys Leu Pro Ser Tyr Ser Thr His Glu Leu Cys Glu Arg Phe  
 1025 1030 1035 1040  
 Ala Arg Ile Met Leu Ser Leu Ser Arg Thr Pro Ala Asp Gly Arg  
 1045 1050 1055

&lt;210&gt; 2

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

Gly Leu Lys Asn Val Gly Asn Thr Cys Trp Phe Ser Ala Val Ile Gln  
 1 5 10 15

Ser Leu

<210> 3  
 <211> 17  
 <212> PRT  
 <213> Homo sapiens

<400> 3  
 Gln Gln Asp Val Ser Glu Phe Thr His Lys Leu Leu Asp Trp Leu Glu  
 1 5 10 15

Asp

<210> 4  
 <211> 21  
 <212> PRT  
 <213> Homo sapiens

<400> 4  
 Tyr Arg Leu His Ala Val Leu Val His Glu Gly Gln Ala Asn Ala Gly  
 1 5 10 15

His Tyr Trp Ala Tyr  
 20

<210> 5  
 <211> 1087  
 <212> PRT  
 <213> Homo sapiens

<400> 5  
 Met Thr Val Glu Gln Asn Val Leu Gln Gln Ser Ala Ala Gln Lys His  
 1 5 10 15

Gln Gln Thr Phe Leu Asn Gln Leu Arg Glu Ile Thr Gly Ile Asn Asp  
 20 25 30

Thr Gln Ile Leu Gln Gln Ala Leu Lys Asp Ser Asn Gly Asn Leu Glu  
 35 40 45

Leu Ala Val Ala Phe Leu Thr Ala Lys Asn Ala Lys Thr Pro Gln Gln  
 50 55 60

Glu Glu Thr Thr Tyr Tyr Gln Thr Ala Leu Pro Gly Asn Asp Arg Tyr

7



325	330	335
Leu Glu Ala Ala Met Ile Glu Gly Glu Ile Glu Ser Leu His Ser Glu		
340	345	350
Asn Ser Gly Lys Ser Gly Gln Glu His Trp Phe Thr Glu Leu Pro Pro		
355	360	365
Val Leu Thr Phe Glu Leu Ser Arg Phe Glu Phe Asn Gln Ala Leu Gly		
370	375	380
Arg Pro Glu Lys Ile His Asn Lys Leu Glu Phe Pro Gln Val Leu Tyr		
385	390	395
Leu Asp Arg Tyr Met His Arg Asn Arg Glu Ile Thr Arg Ile Lys Arg		
405	410	415
Glu Glu Ile Lys Arg Leu Lys Asp Tyr Leu Thr Val Leu Gln Gln Arg		
420	425	430
Leu Glu Arg Tyr Leu Ser Tyr Gly Ser Gly Pro Lys Arg Phe Pro Leu		
435	440	445
Val Asp Val Leu Gln Tyr Ala Leu Glu Phe Ala Ser Ser Lys Pro Val		
450	455	460
Cys Thr Ser Pro Val Asp Asp Ile Asp Ala Ser Ser Pro Pro Ser Gly		
465	470	475
Ser Ile Pro Ser Gln Thr Leu Pro Ser Thr Thr Glu Gln Gln Gly Ala		
485	490	495
Leu Ser Ser Glu Leu Pro Ser Thr Ser Pro Ser Ser Val Ala Ala Ile		
500	505	510
Ser Ser Arg Ser Val Ile His Lys Pro Phe Thr Gln Ser Arg Ile Pro		
515	520	525
Pro Asp Leu Pro Met His Pro Ala Pro Arg His Ile Thr Glu Glu Lys		
530	535	540
Leu Ser Val Leu Glu Ser Cys Leu His Arg Trp Arg Thr Glu Ile Glu		
545	550	555
Asn Asp Thr Arg Asp Leu Gln Glu Ser Ile Ser Arg Ile His Arg Thr		
565	570	575
Ile Glu Leu Met Tyr Ser Asp Lys Ser Met Ile Gln Val Pro Tyr Arg		

580 585 590  
 Leu His Ala Val Leu Val His Glu Gly Gln Ala Asn Ala Gly His Tyr  
 595 600 605  
 Trp Ala Tyr Ile Phe Asp His Arg Glu Ser Arg Trp Met Lys Tyr Asn  
 610 615 620  
 Asp Ile Ala Val Thr Lys Ser Ser Trp Glu Glu Leu Val Arg Asp Ser  
 625 630 635 640  
 Phe Gly Gly Tyr Arg Asn Ala Ser Ala Tyr Cys Leu Met Tyr Ile Asn  
 645 650 655  
 Asp Lys Ala Gln Phe Leu Ile Gln Glu Glu Phe Asn Lys Glu Thr Gly  
 660 665 670  
 Gln Pro Leu Val Gly Ile Glu Thr Leu Pro Pro Asp Leu Arg Asp Phe  
 675 680 685  
 Val Glu Glu Asp Asn Gln Arg Phe Glu Lys Glu Leu Glu Glu Trp Asp  
 690 695 700  
 Ala Gln Leu Ala Gln Lys Ala Leu Gln Glu Lys Leu Leu Ala Ser Gln  
 705 710 715 720  
 Lys Leu Arg Glu Ser Glu Thr Ser Val Thr Thr Ala Gln Ala Ala Gly  
 725 730 735  
 Asp Pro Glu Tyr Leu Glu Gln Pro Ser Arg Ser Asp Phe Ser Lys His  
 740 745 750  
 Leu Lys Glu Glu Thr Ile Gln Ile Ile Thr Lys Ala Ser His Glu His  
 755 760 765  
 Glu Asp Lys Ser Pro Glu Thr Val Leu Gln Ser Ile Met Met Thr Pro  
 770 775 780  
 Asn Met Gln Gly Ile Ile Met Ala Ile Gly Lys Ser Arg Ser Val Tyr  
 785 790 795 800  
 Asp Arg Cys Gly Pro Glu Ala Gly Phe Phe Lys Ala Ile Lys Leu Glu  
 805 810 815  
 Tyr Ala Arg Leu Val Lys Leu Ala Gln Glu Asp Thr Pro Pro Glu Thr  
 820 825 830  
 Asp Tyr Arg Leu His His Val Val Val Tyr Phe Ile Gln Asn Gln Ala

835	840	845
Pro Lys Lys Ile Ile Glu Lys Thr Leu Leu Glu Gln Phe Gly Asp Arg		
850	855	860
Asn Leu Ser Phe Asp Glu Arg Cys His Asn Ile Met Lys Val Ala Gln		
865	870	875 880
Ala Lys Leu Glu Met Ile Lys Pro Glu Glu Val Asn Leu Glu Glu Tyr		
	885	890 895
Glu Glu Trp His Gln Asp Tyr Arg Lys Phe Arg Glu Thr Thr Met Tyr		
	900	905 910
Leu Ile Ile Gly Leu Glu Asn Phe Gln Arg Glu Ser Tyr Ile Asp Ser		
	915	920 925
Leu Leu Phe Leu Ile Cys Ala Tyr Gln Asn Asn Lys Glu Leu Leu Ser		
	930	935 940
Lys Gly Leu Tyr Arg Gly His Asp Glu Glu Leu Ile Ser His Tyr Arg		
945	950	955 960
Arg Glu Cys Leu Leu Lys Leu Asn Glu Gln Ala Ala Glu Leu Phe Glu		
	965	970 975
Ser Gly Glu Asp Arg Glu Val Asn Asn Gly Leu Ile Ile Met Asn Glu		
	980	985 990
Phe Ile Val Pro Phe Leu Pro Leu Leu Leu Val Asp Glu Met Glu Glu		
	995	1000 1005
Lys Asp Ile Leu Ala Val Glu Asp Met Arg Asn Arg Trp Cys Ser Tyr		
1010	1015	1020
Leu Gly Gln Glu Met Glu Pro His Leu Gln Glu Lys Leu Thr Asp Phe		
1025	1030	1035 1040
Leu Pro Lys Leu Leu Asp Cys Ser Met Glu Ile Lys Ser Phe His Glu		
	1045	1050 1055
Pro Pro Lys Leu Pro Ser Tyr Ser Thr His Glu Leu Cys Glu Arg Phe		
	1060	1065 1070
Ala Arg Ile Met Leu Ser Leu Ser Arg Thr Pro Ala Asp Gly Arg		
1075	1080	1085

<210> 6  
 <211> 3803  
 <212> DNA  
 <213> Homo sapiens

<400> 6

```

acagtcggcg tttcgccgcc tgcccgcggt gcccgcgcac gccggccgcc atcgcttcg 60
cgcttggtg gcggggggcgc tgcctccca ggccgtccgc gccgtccct ggagctcggc 120
ggagcgcggc agccagggcc ggccggaggcg cgaggagccg ggccgccaccg ccgccgccgc 180
cgccgccgcc gcggggggcca tgaccgtgga gcagaacgtg ctgcagcaga gcgcggcgca 240
gaagcaccag cagacgtttt tgaatcaact gagagaaatt acgggggatta atgacacca 300
gatactacag caagccttga aggatagtaa tggaaacttg gaattagcag tggctttcct 360
tactgcgaag aatgctaaga cccctcagca ggaggagaca acttactacc aaacagcact 420
tcctggcaat gatagataca tcagtgtggg aagccaagca gatacaaatg tgattgatct 480
cactggagat gataaagatg atcttcagag agcaattgcc ttgagtttgg ccgaatcaaa 540
cagggcattc agggagactg gaataactga tgaggaacaa gccattagca gagttcctga 600
agccagcata gcagagaata aagcatgttt gaagaggaca cctacagaag tttggaggga 660
ttctcgaaac ctttatgata gaaaaagaca ggacaaagct cccgttgggc taaagaatgt 720
tggcaatact tgttggttta gtgctgttat tcagtcatta tttaatcttt tggaaattag 780
aagattagtt ctgaattaca agcctccatc aaatgctcaa gatttacctt gaaacaaaa 840
ggaacatcgg aatttgcctt ttatgcgtga gctgaggtat ctatttgac ttcttgttgg 900
taccaaaagg aagtatgttg atccatcaag agcagttgaa attcttaagg atgctttcaa 960
atcaaatgac tcacagcagc aagatgtgag tgagtttaca cacaaattat tagattggtt 1020
agaagatgcc ttccaaatga aagctgaaga ggagacggat gaagagaagc caaagaacct 1080
catggtagag ttgttctatg gcagattcct ggctgtggga gtacttgaag gtaaaaaatt 1140
tgaaaacact gaaatgtttg gtcagtacct acttcaggtc aatgggttca aagatctgca 1200
tgagtgccta gaagctgcaa tgattgaagg agaaattgag tctttacatt cagagaattc 1260
aggaaaatca ggccaagagc atttggtttac tgaattacca cctgtgttaa catttgaatt 1320
gtcaagattt gaatttaatc aggcattggg aagaccagaa aaaattcaca acaattaga 1380
atttcccaa gttttatatt tggacagata catgcacaga aacagagaaa taacaagaat 1440
taagagggaa gagatcaaga gactgaaaga ttacctcacg gtattacaac aaaggctaga 1500
aagatattta agctatggtt ccggtcccaa acgattcccc ttggtagatg ttcttcagta 1560
tgcattggaa ttgcctcaa gtaaacctgt ttgcacttct cctgttgacg atattgacgc 1620
tagttccccà cctagtgtgt ccataccatc acagacatta ccaagcacia cagaacaaca 1680
gggagcccta tcttcagaac tgccaagcac atcaccttca tcagttgctg ccatttcac 1740
gagatcagta atacacaaac catttactca gtcccggata cctccagatt tgcccatgca 1800
tccggcacca aggcacataa cggaggaaga actttctgtg ctggaaagtt gtttacatcg 1860
ctggaggaca gaaatagaaa atgacaccag agatttgag gaaagcatat ccagaatcca 1920
tcgaacaatt gaattaatgt actctgacaa atctatgata caagttcctt atcgattaca 1980
tgccgtttta gttcaggaag gccaagctaa tgctgggcac tactgggcat atatttttga 2040
tcacgtgaa agcagatgga tgaagtacaa tgatattgct gtgacaaaat catcatgga 2100
agagctagtg agggactctt ttggtgttta tagaaatgcc agtgcatact gtttaatgta 2160
cataaatgat aaggcacagt tcctaataca agaggagttt aataaagaaa ctgggcagcc 2220
ccttgttggg atagaaacat taccaccgga tttgagagat tttgttgagg aagacaacca 2280
acgatttgaa aaagaactag aagaatggga tgcacaactt gccagaaag ctttgcagga 2340
aaagctttta gcgtctcaga aattgagaga gtcagagact tctgtgacaa cagcacaagc 2400
agcaggagac ccagaatatt tagagcagcc atcaagaagt gattttctcaa agcacttgaa 2460

```

```

agaagaaact attcaaataa ttaccaaggc atcacatgag catgaagata aaagtcctga 2520
aacagttttg cagtcggcaa ttaagttgga atatgcaagg ttggttaagt tggcccaaga 2580
agacacccca ccagaaaccg attatcgttt acatcatgta gtggtctact ttatccagaa 2640
ccaggcacca aagaaaatta ttgagaaaac attactagaa caatttggag atagaaattt 2700
gagttttgat gaaaggtgtc acaacataat gaaagttgct caagccaaac tggaaatgat 2760
aaaacctgaa gaagtaaact tggaggaata tgaggagtgg catcaggatt ataggaaatt 2820
cagggaaaca actatgtatc tcataattgg gctagaaaat tttcaaagag aaagttatat 2880
agattccttg ctgttcctca tctgtgctta tcagaataac aaagaactct tgtctaaagg 2940
cttatacaga ggacatgatg aagaattgat atcacattat agaagagaat gtttgctaaa 3000
attaaatgag caagccgcag aactcttcga atctggagag gatcgagaag taaacaatgg 3060
tttgattatc atgaatgagt ttattgtccc atttttgccca ttattactgg tggatgaaat 3120
ggaagaaaag gatatactag ctgtagaaga tatgagaaat cgatgggtgtt cctaccttgg 3180
tcaagaaatg gaaccacacc tccaagaaaa gctgacagat tttttgccaa aactgcttga 3240
ttgttctatg gagattaaaa gtttccatga gccaccgaag ttaccttcat attccacgca 3300
tgaactctgt gagcgatttg cccgaatcat gttgtccctc agtcgaactc ctgctgatgg 3360
aagataaaact gcacactttc cctgaacaca ctgtataaac tctttttagt tcttaaccct 3420
tgccttcctg tcacaggggt tgcttgttgc tgctatagtt ttttaactttt ttttatttta 3480
ataactgcaa aagacaaaat gactatacag actttagtca gactgcagac aataaagctg 3540
aaaatcgcat ggcgctcaga cattttaacc ggaactgatg tataatcaca aatctaattg 3600
attttattat ggcaaaacta tgcttttgcc accttcctgt tgcagtatta ctttgctttt 3660
atcttttctt tctcaacagc ttccattca gtctggatcc ttccatgact acagccattt 3720
aagtgttcag cactgtgtac gatacataat atttggtagc ttgtaaatga aataaagaat 3780
aaagttttat ttatggctac cta 3803

```

&lt;210&gt; 7

&lt;211&gt; 3169

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 7

```

catgaccgtg gagcagaacg tgctgcagca gagcgcggcg cagaagcacc agcagacgtt 60
tttgaatcaa ctgagagaaa ttacggggat taatgacacc cagatactac agcaagcctt 120
gaaggatagt aatggaaact tggaattagc agtggctttc cttactgcga agaattgctaa 180
gaccctcag caggaggaga caacttacta ccaaacagca cttcctggca atgatagata 240
catcagtgtg ggaagccaag cagatacaaa tgtgattgat ctactggag atgataaaga 300
tgatcttcag agagcaattg cttgagttt ggccgaatca aacagggcat tcaggagac 360
tggaataact gatgaggaac aagccattag cagagttctt gaagccagca tagcagagaa 420
taaagcatgt ttgaagagga cacctacaga agtttgagg gattctcgaa acccttatga 480
tagaaaaaga caggacaaag ctcccgttgg gctaaagaat gttggcaata cttgttggtt 540
tagtgctgtt attcagtcatt ttttaattt tttggaattt agaagattag ttctgaatta 600
caagcctcca tcaaatgctc aagatttacc ccgaaaccaa aaggaacatc ggaatttgcc 660
ttttatgctg gagctgaggt atctatttgc acttcttggt ggtacaaaaa ggaagtatgt 720
tgatccatca agagcagttg aaattcttaa ggatgcttcc aaatcaaatg actcacagca 780
gcaagatgtg agtgagttaa cacacaaatt attagattgg ttagaagatg ccttccaaat 840
gaaagctgaa gaggagacgg atgaagagaa gccaaagaac cccatggtag agttgttcta 900
tggcagattc ctggctgtgg gactacttga aggtaaaaaa tttgaaaaca ctgaaatgtt 960
tggtcagtac ccacttcagg tcaatgggtt caaagatctg catgagtgcc tagaagctgc 1020

```

aatgattgaa ggagaaattg agtctttaca ttcagagaat tcaggaaaat caggccaaga 1080  
gcattgggtt actgaattac cacctgtgtt aacatttgaa ttgtcaagat ttgaatttaa 1140  
tcaggcattg ggaagaccag aaaaaattca caacaaatta gaatttcccc aagttttata 1200  
tttgacaga tacatgcaca gaaacagaga aataacaaga attaagaggg aagagatcaa 1260  
gagactgaaa gattacctca cgggtattaca acaaaggcta gaaagatatt taagctatgg 1320  
ttccgggtccc aaacgattcc ccttggtaga tgttcttcag tatgcattgg aatttgctc 1380  
aagtaaacct gtttgactt ctctgttga cgatattgac gctagttccc cacctagtgg 1440  
ttccatacca tcacagacat taccaagcac aacagaacaa caggagccc tatcttcaga 1500  
actgccaagc acatcacctt catcagttgc tgccatttca tcgagatcag taatacacia 1560  
accatttact cagtcccga tacctccaga ttgcccctg catccggcac caaggcacat 1620  
aacggaggaa gaactttctg tgctggaaag ttgtttacat cgctggagga cagaaataga 1680  
aatgacacc agagatttgc aggaagcat atccagaatc catcgaacaa ttgaattaat 1740  
gtactctgac aaatctatga tacaagttcc ttatcgatta catgccgttt tagttcacga 1800  
aggccaagct aatgctggc actactggc atatattttt gatcatcgtg aaagcagatg 1860  
gatgaagtac aatgatattg ctgtgacaaa atcatcatgg gaagagctag tgagggactc 1920  
tttggtggt tatagaaatg ccagtgcata ctgtttaatg tacataaatg ataaggcaca 1980  
gttcctaata caagaggagt ttaataaaga aactgggcag ccccttggtg gtatagaaac 2040  
attaccaccg gatttgagag attttggtga ggaagacaac caacgatttg aaaaagaact 2100  
agaagaatgg gatgcacaac ttgccagaa agctttgcag gaaaagcttt tagcgtctca 2160  
gaaattgaga gagtcagaga cttctgtgac aacagcaca gcagcaggag acccagaata 2220  
tctagagcag ccatcaagaa gtgatttctc aaagcattg aaagaagaaa ctattcaa 2280  
aattaccaag gcatcacatg agcatgaaga taaaagtcct gaaacagttt tgcagtcggc 2340  
aattaagttg gaatatgcaa ggttggttaa gttggcccaa gaagacacc caccagaaac 2400  
cgattatcgt ttacatcatg tagtggctc ctttatccag aaccaggcac caaagaaaat 2460  
tattgagaaa acattactag aacaatttgg agatagaaat ttgagttttg atgaaagggtg 2520  
tcacaacata atgaaagttg ctcaagccaa actggaaatg ataaaacctg aagaagtaaa 2580  
cttgaggaa tatgaggagt ggcacagga ttataggaaa ttcagggaaa caactatgta 2640  
tctcataatt gggctagaaa attttcaaag agaaagttat atagattcct tgctgttcc 2700  
catctgtgct tatcagaata acaaagaact cttgtctaaa ggcttataca gaggacatga 2760  
tgaagaattg atatcacatt atagaagaga atgtttgcta aaattaaatg agcaagccgc 2820  
agaactcttc gaatctggag aggatcgaga agtaacaat ggtttgatta tcatgaatga 2880  
gtttattgtc ccatttttgc cattattact ggtggatgaa atggaagaaa aggatatact 2940  
agctgtagaa gatatgagaa atcgatggtg ttcctacctt ggtcaagaaa tggaaccaca 3000  
cctccaagaa aagctgacag attttttgcc aaaactgctt gattgttcta tggagattaa 3060  
aagtttccat gagccaccga agttacctt atattccacg catgaactct gtgagcgatt 3120  
tgcccgaatc atgttgctcc tcagtcgaac tctgtctgat ggaagataa 3169

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number  
**WO 00/79267 A3**

(51) International Patent Classification<sup>7</sup>: C12N 9/64, A61K 38/48

(21) International Application Number: PCT/GB00/02446

(22) International Filing Date: 22 June 2000 (22.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
9914589.8 22 June 1999 (22.06.1999) GB  
0008161.2 3 April 2000 (03.04.2000) GB

(71) Applicant (for all designated States except US): SCHOOL OF PHARMACY, UNIVERSITY OF LONDON [GB/GB]; 29-39 Brunswick Square, London WC1N 1AX (GB).

(72) Inventor: and

(75) Inventor/Applicants (for US only): NIZETIC, Dean [UK/GB]; School of Pharmacy, University of London, 29-39 Brunswick Square, London WC1N 1AX (GB). GROET, Jürgen [DE/GB]; School of Pharmacy, University of London, 29-39 Brunswick Square, London WC1N 1AX (GB).

(74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

(88) Date of publication of the international search report:  
15 February 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 00/79267 A3

(54) Title: USE OF HTE UBIQUITIN SPECIFIC PROTEASE USP25 IN THE TREATMENT, PROPHYLAXIS AND DIAGNOSIS OF CANCER

(57) Abstract: The product of a gene located at human chromosome 21q11-21 has ubiquitin specific protease activity and binds to ubiquitin, polyubiquitin, ubiquitin-like protein SUMO-3, and proteins which are implicated in the repair/excision of DNA. It is believed to have a role in regulating cell growth, cell growth arrest and/or apoptosis and is implicated in carcinoma growth. The gene has been sequenced and cloned into a microorganism and the product's effect on cell growth is to be investigated. The gene has GenBank accession number AF134213 and has the HUGO-approved name USP25.

# INTERNATIONAL SEARCH REPORT

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C12N9/64 A61K38/48		International Application No PCT/GB 00/02446
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) BIOSIS, EPO-Internal, WPI Data, EMBL, STRAND, EMBASE, CHEM ABS Data		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GROET JUERGEN ET AL: "Bacteria contig map of the 21q11 region associated with Alzheimer's disease and abnormal myelopoiesis in Down syndrome." GENOME RESEARCH, vol. 8, no. 4, April 1998 (1998-04), pages 385-398, XP002152570 ISSN: 1088-9051 cited in the application page 394, left-hand column, line 5 - line 33  <div style="text-align: center;">--- -/--</div>	1-11
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <span style="margin-left: 100px;"><input checked="" type="checkbox"/> Patent family members are listed in annex.</span>		
* Special categories of cited documents: <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*Z* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search  <div style="text-align: center;">27 November 2000</div>		Date of mailing of the international search report  <div style="text-align: center;">13/12/2000</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016		Authorized officer  <div style="text-align: center;">Morawetz, R</div>

Form PCT/ISA/210 (second sheet) (July 1992)



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/02446

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KOHNO TAKASHI ET AL: "Homozygous deletion and frequent allelic loss of the 21q11.1-q21.1 region including the ANA gene in human lung carcinoma." GENES CHROMOSOMES & CANCER, vol. 21, no. 3, March 1998 (1998-03), pages 236-243, XP000943071 ISSN: 1045-2257 cited in the application the whole document	1-11
Y	CAVANI SIMONA ET AL: "Cytogenetic and molecular study of 32 Down syndrome families: Potential leukaemia predisposing role of the most proximal segment of chromosome 21q." BRITISH JOURNAL OF HAEMATOLOGY, vol. 103, no. 1, October 1998 (1998-10), pages 213-216, XP000943387 ISSN: 0007-1048 the whole document	1-11
Y	FUJIWARA TSUTOMU ET AL: "Identification and chromosomal assignment of USP1, a novel gene encoding a human ubiquitin-specific protease." GENOMICS, vol. 54, no. 1, 15 November 1998 (1998-11-15), pages 155-158, XP002153254 ISSN: 0888-7543 the whole document	1-11
Y	DATABASE EMBL 'Online! Accession number AA081200, 29 November 1996 (1996-11-29) HILLIER, L. ET AL.: "Homo sapiens cDNA clone IMAGE:549350 5' similar to WP:K02C4.3 CE01599 UBIQUITIN CARBOXYL-TERMINAL HYDROLASE" XP002152573 the whole document -& HILLIER, L. ET AL.: "Generation and analysis of 280,000 human expressed sequence tags" GENOME RES., vol. 6, no. 9, 1996, pages 807-828, XP002922485 the whole document	1-11

-/--

# INTERNATIONAL SEARCH REPORT

Int. onal Application No  
PCT/GB 00/02446

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DATABASE EMBL 'Online! Accession number AA482201, 24 June 1997 (1997-06-24) STRAUSBERG, R.: "Homo sapiens cDNA clone IMAGE:824710 5' similar to SW:UBP2-YEAST Q01476 UBIQUITIN CARBOXYL-TERMINAL HYDROLASE 2" XP002152574 the whole document	1-11
Y	WILKINSON KEITH D: "Regulation of ubiquitin-dependent processes by deubiquitinating enzymes." FASEB JOURNAL, vol. 11, no. 14, December 1997 (1997-12), pages 1245-1256, XP000943130 ISSN: 0892-6638 cited in the application page 1251, right-hand column, paragraph 4 -page 1252, right-hand column, paragraph 1	1-11
P,X	WO 99 46289 A (HUMAN GENOME SCIENCES INC :NI JIAN (US); ROSEN CRAIG A (US); FERRI 16 September 1999 (1999-09-16) page 38, line 30 -page 41, line 6	1-4,10, 11
P,X	VALERO REBECA ET AL: "USP25, a novel gene encoding a deubiquitinating enzyme, is located in the gene-poor region 21q11.2." GENOMICS, vol. 62, no. 3, 15 December 1999 (1999-12-15), pages 395-405, XP002152572 ISSN: 0888-7543 cited in the application the whole document	1-11
P,X	GROET JURGEN ET AL: "Narrowing of the region of allelic loss in 21q11-21 in squamous non-small cell lung carcinoma and cloning of a novel ubiquitin-specific protease gene from the deleted segment." GENES CHROMOSOMES & CANCER, vol. 27, no. 2, February 2000 (2000-02), pages 153-161, XP000943091 ISSN: 1045-2257 cited in the application the whole document	1-11

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/02446

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9946289 A	16-09-1999	AU 3006799 A	27-09-1999
		EP 1044210 A	18-10-2000
		WO 9931116 A	24-06-1999
<hr/>			

THIS PAGE BLANK (USPTO)